

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 August 2006 (10.08.2006)

PCT

(10) International Publication Number
WO 2006/084033 A1

(51) International Patent Classification:
A61K 36/00 (2006.01)

(21) International Application Number:
PCT/US2006/003686

(22) International Filing Date: 3 February 2006 (03.02.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/649,902 3 February 2005 (03.02.2005) US

(71) Applicant (for all designated States except US): **SIGNUM BIOSCIENCES, INC.** [US/US]; 1 Deer Park Drive, Suite L-2, Monmouth Junction, New Jersey 08852 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **STOCK, Jeffrey** [US/US]; 6 Washington Street, Rocky Hill, New Jersey 08553. **STOCK, Maxwell** [US/US]; 6 Washington Street, Rocky Hill, New Jersey (US). **HAOMING, Gu** [US/US]; 423 Plainsboro Road, Plainsboro, New J 08536 (US). **STOCK, Gregory, B.** [US/US]; 35 Maple Street, Princeton, New Jersey 08542 (US). **LI, Zhu** [US/US]; 216b Halsey Street, Princeton, New Jersey 08540 (US). **WOLANIN, Peter, Michael** [US/US]; 156 Spruce Street, Princeton, New Jersey 08542 (US).

(74) Agent: **LAFEIR, Ann, M.**; LOWENSTEIN SANDLER PC, 65 Livingston Avenue, Roseland, New Jersey 07068 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR ENHANCING COGNITIVE FUNCTION

(57) Abstract: The invention encompasses novel compositions containing a methylation modifying compound isolated from botanical raw materials, including a fruit of the genus *Coffea*, and microbial raw materials, methods of making such compositions, and methods of treating or preventing human disease using such compositions. The methylation modifying compound in botanical extracts of the present invention can dramatically increase the level of methylation of PP2A, thus increasing cognitive function, particularly in persons suffering from or prone to developing Alzheimer's disease.



WO 2006/084033 A1

COMPOSITIONS AND METHODS FOR ENHANCING COGNITIVE FUNCTION

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of priority of U.S. Provisional Application No. 60/649,902 entitled "Enhancing Cognitive Function," filed February 3, 2005. The entire disclosure of this application is incorporated by reference.

GOVERNMENT RIGHTS

[0002] This work is supported at least in part by grants to Dr. Jeffry Stock. The government may have certain rights in this invention.

FIELD OF THE INVENTION

[0003] The present invention relates to treatment or prevention of human disease and to the enhancement of the general health or well-being of a human subject using compositions extracted from herbs and other botanical and microbial materials.

BACKGROUND OF THE INVENTION

[0004] Alzheimer's disease (AD) is one of the most prevalent of progressive brain disorders. Currently an estimated 4.5 million older Americans suffer from AD. According to the latest estimates of the current and future prevalence of AD, the number of older people with AD will grow dramatically as the population ages. Projections indicate that as many as 13 million older Americans will have AD by 2050 unless new ways are found to prevent or treat the disease. AD is currently the third most expensive disease after heart disease and cancer. Treatment and care of those with AD now runs \$100 to \$150 billion a year and costs are projected to rise sharply as the population ages. Currently, pharmacological treatment of AD is primarily based on the use of acetylcholinesterase inhibitors (AChEIs), which have been reported to provide beneficial effects on cognitive, functional, and behavioral symptoms of the disease. Four of the five drugs approved for AD treatment in the U.S. -- donepezil (Aricept®), rivastigmine (Exelon®), galantamine (Reminyl®), and tacrine (Cognex®) -- are AChEIs. The fifth, Memantine (Namenda®), is an N-methyl-D-aspartate ("NMDA") antagonist that alters glutamate signaling. Because only a small fraction of AD patients respond to this type of treatment, any new approach to the treatment or prevention of AD

would have tremendous value. In spite of intensive research, there are no proven preventive agents or agents capable of limiting progression of the disease, and the few used to ameliorate its symptoms have side effects of nausea, vomiting, diarrhea, and even liver damage, yet do not meaningfully slow the disease's underlying course for most patients. AD may well be the most pharmaceutically under-served major disease in the US.

[0005] AD is physically evidenced by amyloid plaques and neurofibrillary tangles in the brain. These pathological markers are associated with cognitive regression and other varied symptoms of the disease. The amyloid plaques, which are the focus of the preponderance of research today on the disease, contain aggregated amyloid β -peptides derived from proteolytic cleavage of the larger amyloid precursor protein. The major component of neurofibrillary tangles is the protein tau, a constituent of the cytoskeleton. Tau is a microtubule-associated protein that functions in brain to regulate the structure and function of axonal microtubules. Over the past decade, several groups have demonstrated that the tau protein found in neurofibrillary tangles is hyperphosphorylated. Tau hyperphosphorylation is thought to destabilize microtubules and thereby contribute to neurodegeneration and the development of AD. Tau hyperphosphorylation results from an imbalance between kinase and phosphatase activities (reviewed in Buee et al., *Brain Res Brain Res Rev* 33(1): 95-130 (2000)). Several serine/threonine protein kinases have been implicated in tau phosphorylation including cyclin-dependent kinase 5 ("cdk-5"), glycogen synthase kinase-3b ("GSK-3b") and MAP kinases. Tau dephosphorylation appears to be primarily mediated by protein phosphatase 2A ("PP2A"). Importantly, recent results suggest that a decrease in PP2A activity is associated with the elevated levels of tau phosphorylation that appear to cause neurofibrillary tangle formation (Planel et al., *J Biol Chem.* 276(36):34298-306 (2001)). Enhancement of this activity therefore may have significant therapeutic value. Since PP2A methylation greatly enhances the formation of a PP2A heterotrimer, it is believed that enhancing PP2A methylation will result in enhanced PP2A activity towards Tau.

[0006] Protein phosphatase 2A (PP2A) associates with a variety of regulatory subunits. (Janssens, V., Gloris, J., *Biochem. J.* 353 (Pt. 3): 417-39 (2001)). The predominant form in neuronal tissue is a trimer composed of a dimeric core composed of a 65 kilodalton (kDa) A subunit and the 36 kDa PP2A catalytic C subunit associated with one of several different regulatory B subunits. Whereas the A and C subunits are present more or less uniformly, the B subunit is variable and confers substrate specificity and subcellular localization to each

PP2A holoenzyme trimer. The number and types of B subunits present is subject to developmental regulation and is cell type specific.

[0007] The variable B subunits of PP2A are classified into four families: (1) the B family with four isoforms (α , β , γ , δ); (2) the B' family with five isoforms (α , β , γ , δ , ϵ); (3) the B'' family; and (4) the B''' family. The PP2A AB_αC heterotrimer appears to be the major phosphatase in brain responsible for dephosphorylation of tau. (Kamibayashi, C. et al., *J. Biol. Chem.* 269 (31): 20139-148 (1994); Sontag, E., et al., *J. Neuropathol. Exp. Neurol.* 63 (4): 287-301 (2004)).

[0008] The alpha-carboxyl of the C terminal leucine residue of the catalytic subunit of PP2A is subject to methyl esterification and methyl-ester hydrolysis, and the methylation state of PP2A regulates heterotrimer formation. (Tokstykh, T. et al., *EMBO J.* 19 (21): 5682-91 (2000); Wu, J. et al., *EMBO J.* 19 (21): 5672-81 (2000); Wei, H. et al., *J. Biol. Chem.* 276 (2): 1570-77 (2001); Yu, XX, et al., *Mol. Biol. Cell* 12 (1): 185-99 (2001)). Two enzymes are involved in controlling the methylation state of PP2A: (1) an S-adenosylmethionine-dependent PP2A-specific protein methyltransferase ("PPMT"), which adds the methyl group and (2) a PP2A-specific protein methylesterase ("PPME"), which removes the methyl group. PP2A methylation promotes PP2A AB_αC trimer assembly. Any deficiency in methylation is expected to preclude PP2A AB_αC heterotrimer formation, thereby leading to a deficiency in tau dephosphorylation, tau hyperphosphorylation and the formation of neurofibrillary tangles. (Vafai, S.B., Stock, J.B., *FEBS Lett.* 518 (1-3): 1-4 (2002)).

[0009] Homocysteine, a sulfur-containing amino acid that can be either remethylated to methionine or undergo a trans-sulfuration reaction to cystathionine, plays a key role in methylation metabolism (see Figure 1). The conversion of homocysteine to methionine occurs in all tissues. Methionine is activated by ATP in the presence of methionine adenosyl transferase (labeled as (1) in Fig. 1) to form the methyl donor, S-adenosylmethionine ("SAM"). SAM-dependent methylation reactions in the presence of SAM-dependent methyltransferases (labeled as (2) in Fig. 1) result in the formation of S-adenosylhomocysteine ("SAH"), which is cleaved by SAH hydrolase (labeled as (3) in Fig. 1) to form adenosine and homocysteine. This reaction is reversible with the equilibrium favoring the condensation of homocysteine and adenosine. Under normal conditions, homocysteine is rapidly methylated, which favors the further cleavage of SAH.

Homocysteine accumulation leads to global decreases in cellular methylation by the condensation of homocysteine with adenosine to form SAH, which acts as a product inhibitor in cellular methylation reactions. In the United States, 5-10 % of the general population has elevated plasma homocysteine, and this imbalance increases to 30-40% in of the elderly. (Selub J., et al., *Ann. Intern. Med.* 131 (5): 331-39 (1999)). The enzymes cystathionine L-synthase; cystathionine Q lyase; betaine homocysteine methyltransferase; and methionine synthase are labeled as (4), (5), (6) and (7), respectively in Fig. 1. See Vafai, S.B., Stock, J.B., *FEBS Lett.* 2: 518 (2002).

[0010] Over the last several years, data has emerged in clinical literature suggesting a direct association between elevated plasma homocysteine and the occurrence of AD. Seshadri et al., (*N Engl J Med* 346 (7): 476-83 (2002)), demonstrated that elevated homocysteine is a risk factor for AD. After adjusting for other AD risk factors, the study concluded that plasma homocysteine levels greater than 14 μ M coincided with about a 2-fold increased risk for developing AD with an additional 40% increased risk with each 5 μ M incremental rise. Other diseases, conditions or disorders associated with elevated plasma homocysteine include, but are not limited to, atherosclerosis; neurodegenerative disorders, such as Parkinson's disease; cerebrovascular disorders (i.e., disorders pertaining to blood vessels in the brain), such as stroke; neuropsychiatric disorders, such as bipolar disorder and schizophrenia; diabetes (type II), and arthritis.

[0011] An analysis of the clinical and basic science literature indicates that a methylation defect resulting from elevated homocysteine could lead to lowered levels of PP2A methylation that would result in lowered PP2A AB_qC, which is believed to lead to tau hyperphosphorylation, neurofibrillary tangle formation, and dementia (Vafai and Stock, *FEBS Lett* 518(1-3): 1-4 (2002)).

[0012] Cellular pathways for removing plasma homocysteine require folate, Vitamin B6 and B12, and thus high homocysteine levels are expected in mice fed diets deficient in these components. This was demonstrated using, male C57BL/J6 mice. One set of 4 week old mice were placed on a diet that contained folate, vitamin B6, and vitamin B12 and another set were fed diets that lacked these vitamins. The mice were allowed free access to both food and water. After nine weeks on their respective diets, each mouse was sacrificed by cervical dislocation. Blood samples were collected for measurement of plasma homocysteine and the brain was removed and quickly frozen in liquid nitrogen for further analysis of tau

phosphorylation. As expected the vitamin-deficient diets caused substantial increases in plasma Hcy and brain SAH. These increases were accompanied by elevated levels of Tau phosphorylation, as shown in Fig. 2. Figure 2 provides Western blots after SDS-PAGE of extracts prepared from the brains of mice raised on normal diets (A and B) and vitamin-deficient diets (C and D). CP13 and PHF1 are monoclonal antibodies that are specific for phosphorylated tau epitopes. TG5 is a monoclonal antibody that recognizes tau independent of its state of phosphorylation; it thereby provides a control showing that total levels of tau expression are unaffected by diet. Mice raised on diets deficient in folate, B12, and B6 had dramatically elevated levels of total plasma homocysteine, brain S-adenosyl homocysteine and elevated levels of tau phosphorylation. S-Adenosyl methionine levels were not significantly affected.

[0013] The demographics of aging in the United States population, combined with a lack of effective treatments, have heightened the need for AD therapies. Moreover, the development of preventives would be an even greater contribution to public health. A protective agent that could be taken over many years to reduce the risk of AD or to substantively delay its onset would be an invaluable breakthrough.

[0014] Coffee has been used for centuries by a diverse range of populations and is presently the most popular beverage worldwide with over 400 billion cups consumed each year. There are many anecdotal reports of the medicinal value of coffee but in spite of its worldwide prevalence, little is really known about its potential medical uses. Some epidemiological studies have suggested an inverse association between coffee consumption and the risk not only of AD, but also of liver cirrhosis, colorectal cancer, cardiovascular mortality, type 2 diabetes and Parkinson's disease. Recent studies have suggested that coffee consumption reduces the risk for AD by as much as 30% (Lindsay et al., *Am J Epidemiol.* 156(5):445-53 (2002)). Various mechanisms for the purported benefits have been suggested, but none have been explored fully enough for these suggestions to be definitive. Moreover, brewed coffee is a complex mixture that contains several pharmacologically active components, including caffeine.

[0015] The present invention provides a definitive mechanism, i.e., enhancement of PP2A methylation, by which to measure benefit to a subject's health and general well-being, and, in particular, cognitive function. An agent that increases the levels of PP2A methylation would help maintain high levels of PP2A ABC heterotrimer formation and may thereby

prevent the demethylation of PP2A and the lowered levels of PP2A ABC heterotrimers that have been associated with tau hyperphosphorylation, neurofibrillary tangle formation, and neurodegeneration in AD.

SUMMARY OF THE INVENTION

[0016] The present invention provides novel compositions that inhibit PP2A demethylation and thereby prevent negative effects of PP2A demethylation associated with diseases such as AD and methods to prepare these compositions. According to one embodiment of the present invention, a composition for promoting general health and well-being in a mammalian subject, including a human, comprises an isolated methylation modifying compound. According to another embodiment, the compound of the composition is isolated from a botanical raw material. According to another embodiment, the compound of the composition is isolated from a microbial raw material. According to another embodiment, the composition inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme. According to another embodiment, the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme. According to another embodiment, the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase. According to another embodiment, the demethylating activity of the protein phosphatase 2A specific protein methylesterase is determined by measuring levels of protein phosphatase 2A methyl esterification. According to another embodiment, the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; coffee; tea; tobacco; and cocoa. According to another embodiment, the botanical raw material comprises a fruit of a species of plant genus *Coffea*. According to another embodiment, the fruit of the plant genus *Coffea* is a coffee bean. According to another embodiment, the botanical raw material comprises a nutritional supplement. According to another embodiment, the nutritional supplement comprises at least one tocopherol. According to another embodiment, the nutritional supplement comprises Vitamin E. According to another embodiment, the botanical raw material comprises a leaf of a tobacco plant. According to another embodiment, the botanical raw material comprises a ground coffee or an instant coffee. According to another

embodiment, the methylation modifying compound is isolated from the botanical raw material by extracting the botanical raw material with a polar solvent. According to another embodiment, the polar solvent is a lower alcohol. According to another embodiment, the lower alcohol is ethanol. According to another embodiment, the polar solvent is ethyl acetate. According to another embodiment, the methylation modifying compound is soluble in ethyl acetate below a pH of about 5 and insoluble in ethyl acetate above a pH of about 10. According to another embodiment, the composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.

[0017] According to yet another embodiment of the present invention, a comestible for promoting general health and well-being in a mammalian subject, including a human, comprises a composition containing an isolated methylation modifying compound. According to another embodiment, the compound is isolated from a botanical raw material. According to another embodiment, the compound is isolated from a microbial raw material. According to another embodiment, the composition inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme. According to another embodiment, the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme. According to another embodiment, the demethylating enzyme is protein phosphatase 2A specific protein methylesterase. According to another embodiment, the level of protein methylesterase demethylating activity is determined by measuring levels of protein phosphatase 2A methyl esterification. According to another embodiment, the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; coffee; tea; tobacco; and cocoa. According to another embodiment, the botanical raw material comprises a fruit of a species of plant genus *Coffea*. According to another embodiment, the fruit of the plant genus *Coffea* is a coffee bean. According to another embodiment, the botanical raw material comprises a nutritional supplement. According to another embodiment, the nutritional supplement comprises at least one tocopherol. According to another embodiment, the nutritional supplement comprises Vitamin E. According to another embodiment, the botanical raw material comprises a leaf of a tobacco plant. According to another embodiment, the botanical raw material comprises a ground coffee. According to another embodiment, the botanical

raw material comprises an instant coffee. According to another embodiment, the methylation modifying compound is isolated from the botanical raw material by extracting the botanical raw material with a polar solvent. According to another embodiment, the polar solvent is a lower alcohol. According to another embodiment, the lower alcohol is ethanol. According to another embodiment, the polar solvent is ethyl acetate. According to another embodiment, the methylation modifying compound is soluble in ethyl acetate below a pH of about 5 and insoluble in ethyl acetate above a pH of about 10. According to another embodiment, the composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid. According to another embodiment, the comestible is a beverage. According to another embodiment, the beverage is selected from the group consisting of a drink comprising water, a fruit drink, a coffee, a tea, an energy drink, a baby formula, an adult nutritional drink, a health drink, and a sports drink. According to another embodiment, the comestible is a food. According to another embodiment, the comestible is a cereal. According to another embodiment, the comestible is a chewing gum. According to another embodiment, the comestible is a candy. According to another embodiment, the comestible is an ingredient of an inhalant. According to another embodiment, the comestible is an ingredient of a transdermal delivery system.

[0018] According to yet another embodiment of the present invention, a pharmaceutical preparation for promoting general health and well-being in a mammalian subject, including a human, comprises a cognitive function-enhancing amount of a composition containing an isolated methylation modifying compound and a pharmaceutically acceptable carrier.

According to another embodiment, the compound is isolated from a botanical raw material. According to another embodiment, the compound is isolated from microbial raw material. According to another embodiment, the compound inhibits a demethylating enzyme that acts on a protein phosphatase 2A and thereby stimulates methylation of the protein phosphatase 2A enzyme. According to another embodiment, the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme. According to another embodiment, the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase. According to another embodiment, the demethylating activity of the protein phosphatase 2A specific protein methylesterase is determined by measuring levels of protein phosphatase 2A methyl esterification. According to another embodiment, the botanical raw material comprises at least one substance selected from the group consisting of ginkgo;

allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline, inositol; tea; tobacco; and cocoa. According to another embodiment, the botanical raw material comprises a fruit of a species of plant genus *Coffea*. According to another embodiment, the fruit of the plant genus *Coffea* is a coffee bean. According to another embodiment, the botanical raw material comprises a ground or instant coffee. According to another embodiment, the botanical raw material is a nutritional supplement. According to another embodiment, the nutritional supplement comprises at least one tocopherol. According to another embodiment, the nutritional supplement comprises vitamin E. According to another embodiment, the botanical raw material comprises a leaf of a tobacco plant. According to another embodiment, the methylation modifying compound is isolated from the botanical raw material by extracting the botanical raw material with a polar solvent. According to another embodiment, the polar solvent is a lower alcohol. According to another embodiment, the lower alcohol is ethanol. According to another embodiment, the polar solvent is ethyl acetate. According to another embodiment, the methylation modifying compound is soluble in ethyl acetate below a pH of about 5 and insoluble in ethyl acetate above a pH of about 10. According to another embodiment, the pharmaceutical preparation is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid. According to another embodiment, the preparation is administered orally. According to another embodiment, the preparation is in the form of a powder. According to another embodiment, the preparation is in the form of a tablet, a capsule, a lozenge, or a suppository.

[0019] According to yet another embodiment of the present invention, a method for promoting general health and well-being in a mammalian subject, including a human, the method comprises the steps (a) isolating a methylation modifying compound; and (b) administering a cognitive function-enhancing amount of a composition comprising the isolated methylation modifying compound to a subject in need thereof. According to another embodiment, in step (a), the methylation modifying compound is isolated from a botanical raw material. According to another embodiment, in step (a) the methylation modifying compound is isolated from a microbial raw material. According to another embodiment, the methylation modifying compound inhibits a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme. According to another embodiment, the composition inhibits at least about 50% of

the demethylating activity of the demethylating enzyme. According to another embodiment, the demethylating enzyme is a protein phosphatase 2A specific protein methyltransferase. According to another embodiment, the method further comprises the step of (c) determining the demethylating activity of the protein phosphatase 2A specific protein methyltransferase. According to another embodiment, the cognition function-enhancing amount of the composition is from about 1 mg to about 10 g. According to another embodiment, the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; tea; tobacco; and cocoa. According to another embodiment, the botanical raw material comprises a fruit of a species of plant genus *Coffea*. According to another embodiment, the fruit of the plant genus *Coffea* is a coffee bean. According to another embodiment, the botanical raw material comprises a ground or instant coffee. According to another embodiment, the botanical raw material comprises a nutritional supplement. According to another embodiment, the nutritional supplement comprises at least one tocopherol. According to another embodiment, the nutritional supplement comprises vitamin E. According to another embodiment, the botanical raw material comprises a leaf of a tobacco plant. According to another embodiment, step (a) further comprises the step of extracting the botanical raw material with a polar solvent. According to another embodiment, the polar solvent is a lower alcohol. According to another embodiment, the lower alcohol is ethanol. According to another embodiment, the polar solvent is ethyl acetate. According to another embodiment, step (a) further comprises the steps of solubilizing the methylation modifying compound in ethyl acetate below a pH of about 5 and partitioning the methylation modifying compound between an aqueous phase and ethyl acetate above a pH of about 10. According to another embodiment, the extracted composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid. According to another embodiment, in step (b), the composition is administered orally. According to another embodiment, the composition is a powder, tablet, capsule, lozenge or suppository.

[0020] According to yet another embodiment of the present invention, a composition for promoting general health and well-being in a mammalian subject, including a human, is prepared by a process, which comprises the steps: a) contacting a botanical raw material with a polar solvent for a time sufficient to form an extract solution from the botanical raw

material; b) isolating the extract solution and removing the polar organic solvent to form a concentrated primary extract; and c) dissolving the composition in a solvent at a pH ranging from about 7 to about 10 to form a concentrated purified extract. According to another embodiment, the polar solvent in step (a) of the process is a lower alcohol. According to another embodiment, the lower alcohol is ethanol. According to another embodiment, the polar solvent in step (a) of the process is ethyl acetate. According to another embodiment, step (a) of the process further comprises the step of solubilizing the composition in ethyl acetate below a pH of about 5. According to another embodiment, step (b) of the process further comprises the step of partitioning the methylation modifying compound between an aqueous phase and ethyl acetate above a pH of about 10. According to another embodiment, the composition inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme. According to another embodiment, the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme. According to another embodiment, the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase. According to another embodiment, the botanical raw material comprises at least one substance selected from the group consisting of ginkgo; allicin; bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng; grape seed; St. John's wort; artichoke-powder; choline; inositol; coffee; tea; tobacco; and cocoa. According to another embodiment, the botanical raw material comprises a fruit of a species of plant genus *Coffea*. According to another embodiment, the fruit of the plant genus *Coffea* is a coffee bean. According to another embodiment, the botanical raw material comprises a ground or instant coffee. According to another embodiment, the botanical raw material comprises a nutritional supplement. According to another embodiment, the nutritional supplement comprises at least one tocopherol. According to another embodiment, the nutritional supplement comprises Vitamin E. According to another embodiment, the botanical raw material comprises a leaf of a tobacco plant. According to another embodiment, the concentrated purified extract is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid. According to another embodiment, the composition is administered orally. According to another embodiment, the composition is in the form of a powder, a tablet, a capsule, a lozenge or a suppository.

[0021] According to yet another embodiment of the present invention, a method of treating or preventing cognitive effects of a disease, condition, or disorder involving defective methylation metabolism in a mammalian subject, including a human, comprises the steps of (a) isolating a methylation modifying compound; and (b) administering a therapeutically effective amount of a composition comprising the isolated methylation modifying compound to a subject in need thereof. According to another embodiment, in step (a) the methylation modifying compound is isolated from a botanical raw material. According to another embodiment, in step (a) the methylation modifying compound is isolated from a microbial raw material. According to another embodiment, the disease, condition, or disorder is associated with higher than normal levels of plasma homocysteine. According to another embodiment, the disease condition or disorder is at least one disease, condition or disorder selected from the group consisting of a cardiac disorder; atherosclerosis; a neurodegenerative disorder; a cerebrovascular disorder; a neuropsychiatric disorder; and diabetes. According to another embodiment, the methylation modifying compound inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme. According to another embodiment, the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme. According to another embodiment, the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase. According to another embodiment, the therapeutically effective amount of the composition is from about 1 mg to about 10 g of the composition. According to another embodiment, the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tacopherol; Vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; tea; tobacco; and cocoa. According to another embodiment, the botanical raw material comprises a fruit of a species of plant genus *Coffea*. According to another embodiment, the fruit of the plant genus *Coffea* is a coffee bean. According to another embodiment, the botanical raw material comprises a ground or instant coffee. According to another embodiment, the botanical raw material comprises a nutritional supplement. According to another embodiment, the nutritional supplement comprises at least one tocopherol. According to another embodiment, the nutritional supplement comprises Vitamin E. According to another embodiment, the botanical raw material comprises a leaf of a tobacco plant. According to another

embodiment, step (a) further comprises the step of extracting the botanical raw material with a polar solvent. According to another embodiment, the polar solvent is a lower alcohol. According to another embodiment, the polar solvent is ethyl acetate. According to another embodiment, step (a) further comprises the steps of solubilizing the methylation modifying compound in ethyl acetate below a pH of about 5 and partitioning the methylation modifying compound between an aqueous phase and ethyl acetate above a pH of about 10. According to another embodiment, the composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid. According to another embodiment, in step (b) the composition is administered orally. According to another embodiment, the composition is a powder, a tablet, a capsule, a lozenge, or a suppository.

Brief Description of the Figures

[0022] Figure 1 provides a schematic diagram of the methyl cycle (1: methionine adenosyl transferase; 2: SAM-dependent methyl transferases; 3: SAH hydrolase; 4: Betaine homocysteine methyl transferases; 5: methionine synthase).

[0023] Figure 2 provides Western blots after SDS-PAGE of extracts prepared from the brains of mice raised on normal diets (A and B) and vitamin-deficient diets (C and D).

[0024] Figure 3 demonstrates that an ethanol extract prepared from coffee grounds shows a concentration dependent inhibition of PP2A methyl esterase.

[0025] Figure 4 parts A through D show purification of PP2A methyl esterase inhibitor compound activity extracted from instant coffee by ethanol extraction: (A) PP2A methyl esterase inhibitor compound activity in crude ethanol extract after three 100 ml ethanol extractions, 300 ml total volume; (B) PP2A methyl esterase inhibitor compound activity in crude ethanol extract after two 100 ml and one 50 ml ethanol extractions, 200 ml total volume; (C) PP2A methyl esterase inhibitor compound activity in water wash compared to inhibitor remaining in the water insoluble material. (D) elution profile from a C18 solid phase extraction cartridge showing further purification of PP2A methyl esterase inhibitor compound.

DETAILED DESCRIPTION OF THE INVENTION

[0026] The present invention encompasses novel compositions extracted from botanical and microbial extracts, methods of making the compositions, and methods of treating or preventing human disease using the novel compositions. The inventive extracts can dramatically decrease the demethylation of PP2A, thus increasing cognitive function and the resilience of cognitive function, particularly in persons suffering from or prone to developing Alzheimer's disease.

[0027] The present invention is based on the recognition that a botanical extract that stimulates methylation of PP2A or inhibits the demethylation of PP2A may be useful to treat certain health problems, including, but not limited to AD, that have been associated with decreases in PP2A methylation. It has long been believed that coffee products, in particular the caffeine in such products, aid in mental focus, learning, and memory. In one embodiment of the present invention, an extract of coffee that is essentially free of caffeine is shown to possess a botanical ingredient having PP2A-specific protein methylesterase inhibitor activity capable of enhancing cognitive function.

Definitions

[0028] The term "active constituent" is defined as the chemical constituent in a botanical raw material or a microbial raw material that is responsible for the intended therapeutic effect.

[0029] "Alkyl" as used herein refers to a straight or branched chain optionally substituted hydrocarbon having from one to 10 carbon atoms. Examples of "alkyl" as used herein include, but are not limited to methyl, ethyl, isopropyl n-butyl, t-butyl, and the like.

[0030] The term "botanical raw material" as used herein refers to a fresh or processed (e.g. cleaned, frozen, dried, sliced, dissolved, or liquefied) part of a single species of plant or a fresh or processed alga or macroscopic fungus.

[0031] The term "botanical ingredient" refers to a component that originates from a botanical raw material.

[0032] The term "botanical product" refers to a finished, labeled product that contains vegetable matter, which may include plant materials, algae, macroscopic fungi, or combinations thereof. Depending in part on its intended use, a botanical product may be a food, drug, medical device or cosmetic.

[0033] The term “botanical extract” as used herein refers to a product prepared by separating, by chemical or physical process, medicinally active portions of a plant from the inactive or inert components. The botanical extracts prepared according to the present invention preferably are obtained by means of a solvent, optionally under pressure and/or heat.

[0034] As used herein, the term “cognitive function” refers to the ability to perform mental tasks, such as thinking, learning, judging, remembering, computing, controlling motor functions, and the like. The expression “resilience of cognitive function” refers to the ability of functional elements of cognitive function to resist deterioration over time. As used herein, the term “cognitive function enhancing amount” refers to that amount of the composition of the present invention that will noticeably impact the ability to perform mental tasks, as measured by tests for memory, computation, attention, or other mental or cognitive attribute, or as suggested by an individual’s perception of his or her abilities in these realms.

[0035] As used herein, the term “comestible” refers to a material that is suitable for human consumption, including a material that can be ingested by oral and by a non-oral means, e.g., an inhalant or a snuff. For purposes of the present invention, the term includes supplemented or enhanced foods.

[0036] The terms “dietary supplement” and “nutritional supplement” are used interchangeably herein to mean (1) a product intended to supplement the diet that bears or contains one or more of the following dietary ingredients: [A] a vitamin, [B] a mineral, [C] an herb or other botanical, [D] an amino acid, [E] a dietary substance for use by man to supplement the diet by increasing the total dietary intake; or (F) a concentrate, metabolite, constituent, extract, or combination of any ingredient described in clause (A), (B), (C), (D), or (E); and (2) a product that (A)(i) is intended for ingestion; (B) is not represented for use as a conventional food or as a sole item of a meal or the diet; and (C) is labeled as a dietary supplement. For purposes of the present invention, this definition includes tobacco.

[0037] The term “essentially free” means less than about 10% of the amount found in unprocessed material. For example, if a coffee bean contains about 1% w/w caffeine, then an extract that is essentially free of caffeine would contain less than about 0.1% caffeine w/w (excluding additional mass due to dilution in water).

[0038] The term “extracting” as used herein refers to the process of drawing out, withdrawing, distilling or otherwise separating one substance from another by a chemical or physical process.

[0039] The term “food” as used herein refers to (a) articles used for food or drink for man or other animals; (2) chewing gum; and (3) articles used for components of any such article.

[0040] The term “health” or “healthy” as used herein refers to a general condition of the body or mind with reference to soundness and vigor, as well as freedom from disease or ailment.

[0041] The term “lower alcohol” refers to a chemical compound of the general form R-OH, where R is an alkyl group having between one and six carbons.

[0042] As used herein, the phrase “methylation modifying compound” refers to an agent that either directly inhibits at least one enzyme that demethylates PP2A, including, but not limited to a PP2A-specific protein methylesterase, or that indirectly affects the methylation state of PP2A itself.

[0043] The term “microbe” or “microorganism” are used interchangeably herein to refer to an organism too small to be seen clearly with the naked eye, including, but not limited to, microscopic bacteria, fungi(molds), algae, protozoa, and viruses.

[0044] A “microbial raw material” as used herein refers to a fresh or processed (e.g. concentrated, frozen, dried, dissolved, liquefied, pelleted) part of a microbial culture.

[0045] The term “microbial ingredient” refers to a component that originates from a microbial raw material.

[0046] The term “microbial product” refers to a finished, labeled product that contains matter derived from a microbial culture.

[0047] The term “partitioning” as used herein refers to a process that takes advantage of the differential solubility of a substance in two solvents.

[0048] The terms “soluble” and “solubility” refer to the property of being susceptible to being dissolved in a specified fluid (solvent). The term “insoluble,” as used herein refers to the property of a material that has minimal or limited solubility in a specified solvent.

[0049] The term “well-being” as used herein refers to a subject’s physical and mental soundness.

Compositions of the Present Invention

[0050] One aspect of the present invention provides compositions comprising a methylation modifying compound isolated from a botanical raw material or microbial raw material (i) that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50%, more preferably by at least 90%, of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification; or (ii) that stimulates the methylating activity of at least one enzyme that methylates PP2A. In a preferred embodiment, the methylation modifying compound comprises the portion of a botanical extract prepared from fruit of a species of the plant genus *Coffea* that is insoluble in ethyl acetate above a pH of about 10, yet that is soluble in ethyl acetate below a pH of about 5. Preferably, the composition is essentially free of caffeine, caffeic acid and/or chlorogenic acid. Caffeine, also known as trimethylxanthine, caffeine, theine, mateine, guaranine, methyltheobromine and 1,3,7-trimethylxanthine, is a xanthine alkaloid found naturally in coffee beans, tea, kola nuts, Yerba mate, guarana berries, and the like. Caffeic acid (3-(3,4-Dihydroxyphenyl)-2-propenoic acid), which is totally unrelated to caffeine, is found in many fruits, vegetables, seasonings and beverages consumed by humans, principally in conjugated forms such as chlorogenic acid. Chlorogenic acid, an ester of caffeic acid and quinic acid, is a major phenolic compound in coffee, isolated from the leaves and fruits of dicotyledonous plants. Chlorogenic acid also slows the release of glucose into the bloodstream after a meal. A skilled artisan will appreciate that if desired, caffeine and/or chlorogenic acid may be added to the composition.

[0051] In one embodiment, the composition of the invention is prepared by extracting coffee beans, the fruit of the coffee tree, either green, roasted or otherwise treated, of *C. arabica*, *C. robusta*, *C. liberica*, *C. arabusta*, or other species. The extraction procedure concentrates or isolates those agents in coffee that increase the methylation levels of PP2A. The composition of the invention includes extracts or compositions of coffee that are selected, isolated, bred, or genetically modified so as to increase the concentrations of the agents or classes of agents having these activities and includes compositions of coffee that are fortified with these agents extracted either from coffee or from other sources.

[0052] Our experiments with coffee extracts indicate that coffee contains agents that prevent the demethylation of PP2A by PPME and that these demethylation inhibitors are distinct from caffeine or its derivatives. Inhibition of this esterase results in an overall higher

level of methyl-PP2A, and this would be expected to prevent tau hyperphosphorylation and tangle formation associated with AD. Our assay for this activity involves incubating the test compound with purified PPME and purified ^3H -methyl-labeled PP2A in 50 mM MOPS buffer at pH 7.2. We have shown that extracts of several herbal substances have a significant inhibitory effect on PPME. We have analyzed a drip-brewed coffee embodiment of the present invention and extracts of coffee embodiments of the present invention in most detail (See Examples 4, 6, and 7). We have discovered that the inhibitor effect is due to a small molecule that is soluble in polar solvents and is not caffeine, chlorogenic acid or caffeic acid.

[0053] In another aspect, the present invention provides comestibles comprising a composition comprising a methylation modifying compound that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50%, more preferably at least 90%, of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification, and wherein the methylation modifying compound is extracted from a botanical or microbial extract. Preferably the composition is extracted from fruit of a species of the plant genus *Coffea*. The comestibles of the invention include fruit-based drinks, coffee-based drinks, tea-based drinks, sport drinks, nutrition bars, snack foods, gums, cereals, candies, baby formulas, energy drinks, adult nutritional drinks, health drinks, and other food products. The term “sports drink” refers to a beverage that is supposed to rehydrate athletes, as well as restoring electrolytes, sugar and other nutrients, for example, Gatorade, POWERade, and All Sport. As used herein, the term “energy drink” refers to a beverage, including, but not limited to, Jolt Cola, Red Bull and similar products, that contains legal stimulants, vitamins and minerals; these products are formulated to give the user a burst of energy. The term “adult nutritional drink” as used herein refers to such products as Ensure, Longetics® or a similar product. The term “health drink” refers to any beverage purported to have beneficial health effects, including, but not limited to, reducing inflammation; supporting the immune system; neutralizing infectious agents; preventing clogged arteries, preserving cognitive function and inhibiting cancer growth. The comestible can comprise an additional ingredient that confers cognitive or other health benefits.

[0054] Yet another aspect of the invention provides pharmaceutical compositions comprising a composition comprising a methylation modifying compound that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50%, more preferably at least 90%, of the demethylating activity of the demethylating enzyme as

measured by levels of PP2A methyl esterification, and wherein the methylation modifying compound is extracted from a botanical or microbial extract. Preferably, the composition is extracted from fruit of a species of the plant genus *Coffea*. The pharmaceutical composition can be formulated for oral consumption or in the form of a suppository. The pharmaceutical compositions of the invention include powders, tablets, capsules and lozenges.

[0055] Another aspect of the invention provides methods of enhancing cognitive function in a human, the method comprising the step of administering a cognitive function enhancing amount of a composition comprising a methylation modifying compound that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50% of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification, and wherein the methylation modifying compound is extracted from a botanical or microbial extract. Preferably the composition is extracted from fruit of a species of the plant genus *Coffea*. Alternatively, a method of enhancing cognitive function in a human according to the present invention comprising the step of administering a comestible comprising a cognitive function enhancing amount of a composition comprising a methylation modifying compound that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50% of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification, and wherein the methylation modifying compound is extracted from a botanical or microbial extract.

[0056] The invention also provides methods of enhancing memory in a human, which method comprises administering a memory enhancing amount of a composition or a comestible comprising methods of enhancing cognitive function in a human, the method comprising the step of administering a cognitive function enhancing amount of a composition comprising a methylation modifying compound that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50% of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification, and wherein the methylation modifying compound is extracted from either a botanical or a microbial extract. According to the present invention, the composition can be used in methods of treating or preventing any disease, condition or disorder where defects in methylation metabolism appear to play a role as evidenced by an association of the disease, condition or disorder with plasma homocysteine levels that are elevated relative to normal plasma homocysteine levels. Such diseases, conditions or disorders include, but are not

limited to, neurodegenerative diseases, disorders or conditions, such as Parkinson's disease, neuropsychiatric diseases, disorders or conditions, such as bipolar disorder, Alzheimer's disease, heart disease, arthritis, diabetes and certain cancers. The term "neurodegenerative" as used herein refers to a disease, condition or disorder marked by the loss or diminution of an original nerve cell function, and the term "neuropsychiatric" relates to organic and functional diseases, conditions or disorders of the nervous system.

[0057] Yet another aspect of the invention provides processes for preparing a composition comprising a methylation modifying compound that inhibits at least one enzyme that demethylates PP2A, wherein the composition inhibits at least 50%, more preferably at least 90%, of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification, and wherein the methylation modifying compound is extracted from either a botanical extract or a microbial extract. When the composition of the present invention is extracted from the fruit of a species of the plant genus *Coffea*, the process of the present invention comprises the steps: determining when the composition inhibits at least 50%, more preferably at least 90%, of the demethylating activity of the demethylating enzyme as measured by levels of PP2A methyl esterification, and optionally treating the fruit to remove caffeine. The extract can be treated further to remove chlorogenic acid and/or caffeic acid.

Methods of Preparing Compositions of the Invention

[0058] Another aspect of the invention provides a method of preparing a concentrated purified extract comprising a composition useful for enhancing or maintaining cognitive health. One method of preparing a composition of the invention is to extract the active agents from botanical raw material or microbial raw material into organic solvents from aqueous solutions at acid or neutral pH. Further concentration of the agents can be effected by extracting them from organic solvents back into aqueous solvents at basic pH. In one preferred embodiment involving a botanical raw material, the method comprises the steps : (a) contacting the fruit of a species of plant with a polar organic solvent for a time sufficient to form an extract solution, b) removing particulate matter from the extract solution; (c) isolating the extract solution and removing the polar organic solvent to form a concentrated primary extract, (d) washing the concentrated primary extract with a solvent in which impurities are soluble and the agent is poorly soluble; (e) removing the solvent; and (f)

dissolving the desired methylation modifying agent in a polar organic solvent or in water at neutral to basic pH. For example, a coffee extract is prepared by:

[0059] a) contacting a species of the fruit of a species *Coffea* with pure ethanol at an elevated temperature (80 °C) for about 5-10 minutes, i.e., a time sufficient to form an ethanol extract solution from the fruit.

[0060] b) removing particulate matter from the ethanol extract solution by filtration or centrifugation;

[0061] c) isolating the ethanol extract solution and removing the ethanol by evaporation in a glass flask to form a concentrated primary extract;

[0062] d) washing the concentrated extract with deionized water at a pH below about 5;

[0063] e) drying the extract to remove residual water; and

[0064] f) dissolving the extract in ethanol to form a washed concentrated primary extract.

[0065] The term “solvent” as used herein refers to a substance, usually liquid, capable of dissolving or dispersing one or more other substances. Chemists have classified solvents into two broad categories according to their polarity: polar and nonpolar. A common measure of the polarity of a solvent is the dielectric constant. The term “polar solvent” as used herein refers to a compound that is composed of polar molecules. A “polar molecule” is one in which there is some separation of charge in the chemical bonds, so that one part of the molecule has a slight positive charge and the other a slight negative charge. Polar solvents may be further classified as protic or aprotic. The term “protic” refers to a hydrogen atom attached to an electronegative atom, while the term “aprotic” refers to a molecule that does not contain an O-H bond. A “polar protic solvent” can be represented by the general formula ROH; the polarity of the polar protic solvent stems from the bond dipole of the O-H bond. Examples of polar protic solvents include, but are not limited to, water, alcohols, and acetic acid. A “dipolar aprotic solvent” is one that contains a bond that has a large bond dipole. Typically, this bond is a multiple bond between carbon and either oxygen or nitrogen. Most dipolar aprotic solvents contain a C=O double bond. Examples of dipolar aprotic solvents include, but are not limited to, acetone and ethyl acetate. As the number of -CH₂- groups in ROH increases, and the relative amount of hydrocarbon character increases, the polarity decreases. The term “nonpolar solvent” refers to compounds that have low dielectric

constants and are not miscible with water. Examples of nonpolar solvents include, but are not limited to benzene, carbon tetrachloride, and diethyl ether.

[0066] Optionally, the washed concentrated primary extract can be dissolved in water and the pH of the washed concentrated primary extract neutralized. The extract then can be further purified by utilizing the property that the desired agent is insoluble in ethyl acetate above a pH of about 10.0, and is soluble in ethyl acetate below a pH of about 5.0. It would be apparent to a skilled artisan that by assaying for the desired effects on PP2A methylation (see PCT/US03/07658; the contents of which are incorporated by reference) one can readily identify growth conditions and plant varieties having increased levels of the agents of the compositions of the present invention. Likewise, using methods known in the art, a skilled artisan could genetically engineer or breed plant varieties to express increased amounts of the desired agents.

[0067] The compositions of the present invention may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, solutions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs, pastes, gels or the like. Compositions intended for oral use may be prepared according to any known method, and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents, and preserving agents in order to provide pharmaceutically elegant and palatable compositions. Tablets may contain the active ingredient(s) in admixture with non-toxic pharmaceutically-acceptable excipients which are suitable for the manufacture of tablets. These excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch or alginic acid; binding agents, for example, starch, gelatin or acacia; and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They also may be coated for controlled delivery. For example, a "delayed release" dosage form releases a product or substance at a time other than promptly after administration. Examples of delayed-release systems include repeat-action tablets and capsules, and enteric-coated tablets where timed release is achieved by a barrier coating.

[0068] Compositions of the present invention also may be formulated for oral use as hard gelatin capsules, where the active ingredient(s) is(are) mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or soft gelatin capsules wherein the active ingredient(s) is (are) mixed with water or an oil medium, for example, peanut oil, liquid paraffin, or olive oil.

[0069] The compositions of the present invention may be formulated as aqueous suspensions wherein the active ingredient(s) is (are) in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth, and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide such as lecithin, or condensation products of an alkylene oxide with fatty acids, for example, polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example, heptadecaethyl-eneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions also may contain one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[0070] Compositions of the present invention may be formulated as oily suspensions by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil, such as liquid paraffin. The oily suspensions may contain a thickening agent, for example, beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral composition. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

[0071] Compositions of the present invention may be formulated in the form of dispersible powders and granules suitable for composition of an aqueous suspension by the addition of water. The active ingredient in such powders and granules is provided in admixture with a dispersing or wetting agent, suspending agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified

by those already mentioned above. Additional excipients, or example, sweetening, flavoring and coloring agents also may be present.

[0072] The compositions of the invention also may be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oil, or a mineral oil, for example a liquid paraffin, or a mixture thereof. Suitable emulsifying agents may be naturally-occurring gums, for example, gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate. The emulsions also may contain sweetening and flavoring agents.

[0073] The compositions of the invention also may be formulated as syrups and elixirs. Syrups and elixirs may be formulated with sweetening agents, for example, glycerol, propylene glycol, sorbitol or sucrose. Such formulations also may contain a demulcent, a preservative, and flavoring and coloring agents. Demulcents are protective agents employed primarily to alleviate irritation, particularly mucous membranes or abraded tissues. A number of chemical substances possess demulcent properties. These substances include the alginates, mucilages, gums, dextrans, starches, certain sugars, and polymeric polyhydric glycols. Others include acacia, agar, benzoin, carbomer, gelatin, glycerin, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, propylene glycol, sodium alginate, tragacanth, hydrogels and the like.

[0074] The compositions of the present invention also may be in the form of a sterile injectable aqueous or oleaginous suspension. Injectable compositions, such as sterile injectable aqueous or oleaginous suspensions, may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable composition may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1, 3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For parenteral application, "parenteral" meaning subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants. Aqueous suspensions may contain

substances which increase the viscosity of the suspension and include, for example, sodium carboxymethyl cellulose, sorbitol and/or dextran. Optionally, the suspension may also contain stabilizers.

[0075] The terms “drug carrier”, “carrier”, or “vehicle” are used interchangeably to refer to carrier materials suitable for administration of the methylation modifying compounds. Carriers and vehicles useful herein include any such materials known in the art which are nontoxic and do not interact with other components. As used herein the term “a pharmaceutically acceptable carrier” refers to any substantially non-toxic carrier conventionally useable for administration in which the compound will remain stable and bioavailable.

[0076] The compositions of the present invention may be in the form of suppositories for rectal administration of the composition. These compositions can be prepared by mixing the active ingredient with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug. When formulated as a suppository the compositions of the invention may be formulated with traditional binders and carriers, such as triglycerides.

[0077] Compositions of the present invention optionally can include an excipient. The term “excipients” as used herein refers to pharmaceutically acceptable organic or inorganic carrier substances which do not deleteriously react with the active compounds. Suitable dietary excipients include, but are not limited to, dietary suitable starch, vegetable oil, vegetable gums, gelatins, soy extracts, sugars, grains, natural and artificial flavorings, and the like. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil; fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethylcellulose, polyvinylpyrrolidone, and the like. Suitable excipients are described in detail in Remington’s Pharmaceutical Sciences, Twentieth Edition, © 2000 incorporated herein by reference.

[0078] Pharmaceutical compositions can be sterilized and if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds.

[0079] The therapeutically active agent of the present invention can be formulated per se or in salt form. The term “pharmaceutically acceptable salts” refers to nontoxic salts of the active agent. Pharmaceutically acceptable salts include, but are not limited to, those formed with free amino groups such as those derived from hydrochloric, phosphoric, sulfuric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

[0080] Additional compositions of the present invention can be readily prepared using technology which is known in the art such as described in Remington’s Pharmaceutical Sciences.

[0081] A composition of the present invention, alone or in combination with other active ingredients, may be administered to a subject in a single dose or multiple doses over a period of time, generally by oral administration. As used herein, the terms “therapeutically effective amount,” “memory-enhancing amount”, and “cognition enhancing amount” are used interchangeably to refer to the amount of the composition of the invention that results in a therapeutic or beneficial effect, including a subject’s perception of health or general well-being, following its administration to a subject.

[0082] It is believed that an increase in the level of PP2A methylation will bring about the protection or enhancement of cognitive functioning, or preventing a cognitive disorder from manifesting or deepening. Thus the therapeutic effect of the compositions of the present invention can exert a protective or enhancing effect on cognitive function; minimize, prevent or ameliorate cognitive symptoms of a disease or disorder, or may have any other beneficial effect.

[0083] The concentration of the substance is selected so as to exert its therapeutic effect, but low enough to avoid significant side effects within the scope and sound judgment of the skilled artisan. The effective amount of the composition may vary with the age and physical condition of the biological subject being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the specific compound, composition or other active ingredient employed, the particular carrier utilized, and like factors. Those of skill in the art can readily evaluate such factors and, based on this information, determine the particular effective concentration of a composition of the present invention to be used for an intended purpose.

[0084] A skilled artisan can determine a therapeutically effective amount of the inventive compositions by determining the unit dose. As used herein, a “unit dose” refers to the amount of inventive composition required to produce a response of 50% of maximal effect (i.e. ED₅₀). The unit dose can be assessed by extrapolating from dose-response curves derived from in vitro or animal model test systems. The amount of compounds in the compositions of the present invention which will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. (See, for example, Goodman and Gilman’s THE PHARMACOLOGICAL BASIS OF THERAPEUTICS, Joel G. Harman, Lee E. Limbird, Eds.; McGraw Hill, New York, 2001; THE PHYSICIAN’S DESK REFERENCE, Medical Economics Company, Inc., Oradell, N.J., 1995; and DRUG FACTS AND COMPARISONS, FACTS AND COMPARISONS, INC., St. Louis, Mo., 1993). The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient’s circumstances. Various administration patterns will be apparent to those skilled in the art.

[0085] The dosage ranges for the administration of the compositions of the present invention are those large enough to produce the desired therapeutic effect. Preferably, the cognitive function enhancing amount of the compositions of the present invention is administered one or more times per day on a regular basis. A typical dose administered to a human is between about 1 mg and about 10 g of the composition, preferably between 1 mg and 1 g of the composition.

[0086] Those skilled in the art will recognize that initial indications of the appropriate therapeutic dosage of the compositions of the invention can be determined in in vitro and in vivo animal model systems, and in human clinical trials. One of skill in the art would know to use animal studies and human experience to identify a dosage that can safely be administered without generating toxicity or other side effects. For acute treatment where it is desirable to substantially increase methylated PP2A, it is preferred that the therapeutic dosage be close to the maximum tolerated dose. For chronic preventive use, lower dosages may be desirable because of concerns about long term effects. However, coffee and coffee extracts are commonly believed to be safe and have a history of human use. The composition preferably is administered one or more times per day, in food or beverage, as inhalant, in a

gum, transdermally, as a suppository, or as a snuff. As used herein, the term “inhalants” refers to substances or combinations of substances that, by virtue of their high vapor pressure, can be carried by an air current into the nasal passage where they exert their effect. The container from which the inhalant is administered is “an inhaler.”

[0087] Alternatively, the composition of the present invention may be administered at least once per day in combination with a prescribed drug. For example, the composition of the present invention may be administered together with existing anti-cholinesterase drugs now prescribed for Alzheimer's, with various anti-inflammatory agents, or with statins.

[0088] In another aspect, the composition of the present invention is administered at least once per day in combination with a dietary or nutritional supplement believed to have beneficial health effects. For example,

[0089] Coenzyme Q₁₀ (also known as CoQ₁₀, Q₁₀, vitamin Q₁₀, ubiquinone and ubiquinone), a benzoquinone compound synthesized naturally by the human body, is used by cells of the body in oxidative metabolism or cell respiration and as an endogenous antioxidant. An “antioxidant” is a substance that protects cells from free radicals, which are highly reactive chemicals often containing oxygen atoms, that are capable of damaging important cellular components, such as DNA and lipids. The plasma level of CoQ₁₀ has been used in studies as a measure of oxidative stress, a situation in which normal antioxidant levels are reduced. Various investigations have explored the usefulness of CoQ₁₀ as a treatment for diseases, including, but not limited to, cancer and cardiovascular disease.

[0090] Idebenone, a synthetic analog of CoQ₁₀, has been investigated in elderly patients with dementia. Studies suggest that it may diminish nerve cell damage due to ischemia and facilitate memory and learning.

[0091] Huperzine A, a natural acetylcholinesterase inhibitor derived from the Chinese herb *Huperzia serrata*, has antioxidant and neuroprotective properties, and has been proposed as a disease-modifying treatment for AD.

[0092] Galantamine, an acetylcholinesterase inhibitor, is used to treat symptoms of AD.

[0093] Vincamine and vinpocetine, a semisynthetic derivative of vincamine, an alkaloid derived from the plant *Vina minor L*, are used in Europe, Japan and Mexico as pharmaceutical agents for the treatment of cerebrovascular and cognitive disorders.

[0094] Acetyl-L-carnitine, an acetylated derivative of carnitine, has been shown to promote fatty acid beta-oxidation in liver and to prevent motor nerve condition velocity slowing in diabetic rats.

[0095] Dehydroepiandrosterone (DHEA), a steroid, is being studied in the prevention of cancer. In the body, it is a precursor produced by the adrenal gland and converted to testosterone or the estrogens.

[0096] Phosphatidylcholine, a phospholipid that is a major component of cell membranes, has putative activity as a cognition enhancer and in cell-membrane repair

[0097] Gingko, an herb, has putative properties as a neuroprotective agent, an antioxidant, a free-radical scavenger, a membrane stabilizer, and an inhibitor of platelet-activating factor. Sherpina, V.S., et al., *American Family Physician* 68(5) 923-926 (2003). Gingko extract also has been shown to inhibit beta-amyloid deposition. *Id.*

[0098] Curcumin, an active ingredient in turmeric, which is in curry, purportedly has anti-inflammatory and cholesterol lowering properties.

[0099] Ginseng, a Chinese herb, has been used for centuries in Asia as a cure for many maladies.

[00100] Research has shown that Vitamin E (DL-alpha-tocopherol), an essential vitamin that functions as an antioxidant, can help prevent cardiovascular disease and increase the immune response. It has been hypothesized that Vitamin E and its analogs and derivatives may prevent brain cell damage by destroying toxic free radicals. The term "tocol" generally refers to 2-methyl-2-(4,8,12-trimethyltridecyl)chroman-6-ol; the term "tocopherol" generally refers to all mono, di, and trimethyltocols, including, but not limited to, alpha-tocopherol (5,7,8-trimethyltolcol), beta-tocopherol (5,8-dimethyltolcol), gamma-tocopherol (7,8-dimethyltolcol), delta-tocopherol (8-methyltolcol), the term "tocotrienol" refers to 2-methyl-2-(4,8,12-trimethyltrideca-3,7,11-trienyl)chroman-6-ol; and the term "vitamin E" generally refers to all tocol and tocotrienol derivatives exhibiting qualitatively the biological activity of alpha-tocopherol.

[00101] It is well-known that N-acetyl-cysteine (NAC) promotes cellular glutathione production, and thus reduces, or even prevents, oxidant mediated damage. Treatment with NAC provides beneficial effects in a number of respiratory, cardiovascular, endocrine, infectious, and other disease settings.

[00102] B vitamins, such as folic acid, are known to reduce levels of homocysteine, an amino acid already linked, at high levels, to an increased risk of heart attacks, strokes and Alzheimer's disease.

[00103] Lecithin, a lipid material composed of choline and inositol, is a major component of cell membranes. As used by producers of lecithin for commercial use, the term "lecithin" refers to a complex mix of phosphatides and other substances that contain phosphatidylcholine.

[00104] Choline (trimethyl ethanolamine), a quaternary saturated amine classified as an essential nutrient by the Food and Nutrition Board of the Institute of Medicine, is a component of lecithin. Choline is needed by the body to make the neurotransmitter acetylcholine.

[00105] Fish oil, which is oil derived from the tissues of oily fish, naturally contains the omega-3 fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Some experts believe that fish oil can help regulate cholesterol in the body. It also may help protect the brain from cognitive problems associated with Alzheimer's disease.

[00106] Deprenyl (selegiline, Eldepryl®), a monoamine oxidase inhibitor, is prescribed for the treatment of early-stage Parkinson's disease and senile dementia.

[00107] The compositions of the invention can be used alone or in combination with other pharmaceuticals or herbals to prolong mental health, to maintain or enhance cognitive functioning or memory, or to preserve mental or physical well-being and health. The compositions can also be used to prevent or treat effects a number of ailments, including, but not limited to, Alzheimer's disease; Parkinson's disease; heart disease; arthritis; age-related degeneration, functional impairments, and diseases; diabetes, and cancer, have on cognitive function.

[00108] The effectiveness of the compositions and methods of the present invention can be assayed by a variety of protocols. The effects of increasing cognitive function in a human subject can be determined by methods routine to those skilled in the art including, but not limited to, both paper and pencil, and computer tests. One of skill in the art can also directly measure PP2A methylation levels, tau protein phosphorylation levels, neurofibrillary tangle formation and neurodegeneration in animal models.

[00109] Comestibles of the Invention

[00110] Compositions of the present invention may be included in a variety of forms, including, but not limited to, nutritional supplements, pharmaceutical compositions, vitamin supplements, food additives or foods supplements. For example, the methylation modifying compounds described may be embodied as a lozenge, candy, drink, gum, or other snack, food, pill, snuff, nutritional or food supplement, or delivered as an ingredient of an inhalant, beverage, or transdermal delivery system. For example, the methylation modifying compounds of the present invention may be added directly to a liquid beverage, such as water.

[00111] The compositions of the present invention may be in the form of a dispersible dry powder for pulmonary delivery. Dry powder compositions may be prepared by processes known in the art, such as lyophilization and jet milling, as disclosed in International Patent Publication No. WO 91/16038 and as disclosed in U.S. Pat. No. 6,921,527, the disclosures of which are incorporated by reference. The composition of the present invention is placed within a suitable dosage receptacle in an amount sufficient to provide a subject with a unit dosage treatment. The dosage receptacle is one that fits within a suitable inhalation device to allow for the aerosolization of the dry powder composition by dispersion into a gas stream to form an aerosol and then capturing the aerosol so produced in a chamber having a mouthpiece attached for subsequent inhalation by a subject in need of treatment. Such a dosage receptacle includes any container enclosing the composition known in the art such as gelatin or plastic capsules with a removable portion that allows a stream of gas (e.g., air) to be directed into the container to disperse the dry powder composition. Such containers are exemplified by those shown in U.S. Pat. Nos. 4,227,522; U.S. Pat. No. 4,192,309; and U.S. Pat. No. 4,105,027. Suitable containers also include those used in conjunction with Glaxo's Ventolin® Rotohaler brand powder inhaler or Fison's Spinhaler® brand powder inhaler. Another suitable unit-dose container which provides a superior moisture barrier is formed from an aluminum foil plastic laminate. The pharmaceutical-based powder is filled by weight or by volume into the depression in the formable foil and hermetically sealed with a covering foil-plastic laminate. Such a container for use with a powder inhalation device is described in U.S. Pat. No. 4,778,054 and is used with Glaxo's Diskhaler® (U.S. Pat. Nos. 4,627,432; 4,811,731; and 5,035,237). All of these references are incorporated herein by reference.

[00112] The terms “transdermal delivery system”, “transdermal patch”, or “patch” refer to an adhesive system placed on the skin to deliver a time released dose of a drug(s) by passage

from the dosage form through the skin to be available for distribution via the systemic circulation. Transdermal patches are a well-accepted technology used to deliver a wide variety of pharmaceuticals, including, but not limited to, scopolamine for motion sickness, nitroglycerin for treatment of angina pectoris, clonidine for hypertension, estradiol for post-menopausal indications, and nicotine for smoking cessation. Patches suitable for use in the present invention include, but are not limited to, (1) the matrix patch; (2) the reservoir patch; (3) the multi-laminate drug-in-adhesive patch; and (4) the monolithic drug-in-adhesive patch; TRANSDERMAL AND TOPICAL DRUG DELIVERY SYSTEMS, pp. 249-297 (Tapash K. Ghosh *et al.* eds., 1997), hereby incorporated herein by reference. These patches are well known in the art and generally available commercially.

[00113] In preferred embodiments, the comestibles of the invention include a coffee without caffeine, a decaffeinated coffee, a coffee-like beverage, a ready-to-drink coffee, a coffee dessert or any other coffee product enriched for, or selected for, PP2A methylation activity and used for the health functions described above.

[00114] When provided to non-human mammalian subjects, the compositions of the present invention may be administered separately or may be combined with ordinary feed or liquid nourishment to effect the alterations in body composition as described herein.

[00115] The compositions of the invention can be combined with ordinary foods to enhance the value of the palatability of the composition. For example, the compositions can be mixed with soft drinks, food supplements, candy, or high-energy bars, and virtually any other food that can be supplemented with a powder or liquid. Thus, the invention specifically includes food substances of specific types combined with the composition of the invention in specified forms and quantities.

[00116] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges which may independently be included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also included in the invention.

[00117] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

EXAMPLES

[00118] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (*e.g.* amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1: Composition of pure PP2A Methyltransferase

[00119] This example provides a method of preparing pure PPMT for use in assays of demethylation of PP2A. PPMT was obtained using a bacterial overexpression system consisting of the pMT and pBADESL plasmids in *Escherichia coli* strain C41(DE3) (obtained from S. Djordjevic, University College London) (George, R.R. et al., Protein Expr. Purif. 26(2): 266-74 (2002). LB flasks containing 0.2% (w/v) arabinose were inoculated and grown at 37°C to a density of $A_{600} = 0.5$. 1 mM isopropyl-b-D-thiogalactoside (IPTG), an artificial inducer of the Lac operon, was added to induce methyltransferase expression and after 3 hr the culture was centrifuged and the pellet washed with 0.1 M potassium phosphate, pH 7. The pellet was resuspended in 50 mM Tris, pH 7, containing 300 mM NaCl, 10% (v/v) glycerol, 10 mM 2-mercaptoethanol, sonicated on an ice/NaCl bath, and centrifuged. The supernatant was loaded on a pre-equilibrated 10 ml Ni-NTA superflow column, and eluted with a 150 ml gradient from 0 to 300 mM imidazole. The methyltransferase containing fractions were collected, pooled, concentrated and loaded onto a Sephacryl S-100 gel

filtration column, eluted, and fractions collected. The presence of PPMT in the fractions was confirmed by SDS–PAGE analysis. Relevant fractions were pooled and loaded onto a pre-equilibrated Toyopearl Super-Q 650S column. The column then was washed, and the proteins eluted with a gradient of 0 to 500 mM NaCl. Fractions containing pure PPMT were pooled, dialyzed against 50 mM 4-morpholinopropanesulfonic acid, 3-(N-Morpholino)-propanesulfonic acid (MOPS) buffer containing 10% (v/v) glycerol, 1 mM dithiothreitol (DTT), 1 mM ethylenediaminetetraacetic acid (EDTA), and stored at -20°C with 40% glycerol. The protocol yielded approximately 10 mg of pure PPMT per liter of culture media.

Example 2: Pure PP2A Methylesterase

[00120] This example provides a method of preparing pure PPME for use in assays of PP2A methyl esterification. A PP2A methylesterase plasmid (obtained from S. Djordjevic, University College London) was transformed into *E. coli* strain BL21(DE3) cells. LB flasks were inoculated with cultures grown overnight and allowed to grow at 37°C. When the culture reached a density of $A_{600} = 0.5$, 1 mM IPTG was added to induce esterase expression. After 3 hr the cultures were centrifuged and the pellet was washed with 0.1 M potassium phosphate, pH 7, and resuspended in 50 mM Tris buffer containing 300 mM NaCl, 10% glycerol, 10 mM 2-mercaptoethanol. Cells were then disrupted by sonication and the supernatant was loaded on a pre-equilibrated 10 ml Ni: NTA superflow column. The column was washed with Tris-buffer and proteins were eluted with a 150 ml gradient from 0 to 300 mM imidazole. The methylesterase containing fractions were collected, pooled and dialyzed against 50 mM Tris buffer containing 5 % (w/v) glycerol and 1 mM DTT. The proteins were then loaded onto a pre-equilibrated Toyopearl Super-Q 650S column. The column was washed and proteins collected by applying a gradient from 0 to 500 mM NaCl. Fractions containing pure PPME were pooled, dialyzed against 50 mM MOPS buffer containing 10% (v/v) glycerol, 1 mM DTT, 1 mM EDTA and stored at -20°C with 40% glycerol. This protocol gives approximately 10 mg of pure PPME per liter of culture media.

Example 3: Purification of PP2A AC Dimers

[00121] This example provides a method of preparing pure PP2A AC dimers for use in assays of PP2A methyl esterification. (Lee, J., Stock, J., J. Biol. Chem. 268 (26) 19192-195 (1993); Tolstykh, T. et al., EMBO J. 19 (21): 5682-91 (2000)). Two frozen bovine brains were suspended into 800 ml of Buffer A (20 mM MOPS-Na, pH 7.2, 1.0 mM EDTA/DTT and 0.5 mg/L of aprotinin, leupeptin and pepstatin (“protease inhibitors”), and blended until

smooth at 4 °C. 25% $(\text{NH}_4)_2\text{SO}_4$ was added to the brains, stirred for 30 min and centrifuged. $(\text{NH}_4)_2\text{SO}_4$ was added to the supernatant to 70%, stirred for 60 min, and centrifuged. The precipitate in the pellet was collected and dissolved in 200 ml Buffer A and then dialyzed at 4 °C against three, 4 liter changes of Buffer A. The dialyzed solution was clarified by centrifugation and the supernatant was loaded onto a DEAE-Toyopearl 650M column pre-equilibrated with Buffer A. The column was washed and PP2A eluted with Buffer A + 0.3 M NaCl. The eluted PP2A was concentrated in 60 ml Buffer B (50 mM MOPS, pH 7.2, 1 mM EDTA/DTT, 0.5 mg/ml protease inhibitors) with 0.8M $(\text{NH}_4)_2\text{SO}_4$, loaded onto a TSK phenyl column and washed in this buffer before elution with a 1.0 L linear gradient from 0.8M $(\text{NH}_4)_2\text{SO}_4$ to 20% ethylene glycol in Buffer B. Fractions with phosphatase activity were collected and concentrated with a Centriprep concentrator before being loaded on a Sephacryl S-200 column pre-equilibrated in Buffer B with 0.2M NaCl. Fractions with phosphatase activity were pooled, buffer exchanged, and concentrated to 2 ml in Buffer C (50 mM MOPS, pH 7.2, 1 mM EDTA, 1 mM DTT) using a Centriprep concentrator. Concentrated protein was loaded on a Source-15Q HPLC column. A gradient from 0.2M to 0.35 M NaCl in Buffer C in 30 minutes, then from 0.35 M NaCl to 0.5 M NaCl in Buffer C in 10 minutes was used to elute the PP2A. Fractions containing the pure PP2A were collected, concentrated using a Centriprep concentrator, and stored at -20°C with 40% glycerol.

Example 4: Assay for Methylesterase Inhibition

[00122] This example provides a method by which the rate of demethylation of methyl PP2A can be determined. For the composition of ^3H -labeled methyl-PP2A, 100 μl reaction mixtures consisting of purified PP2A AC dimers (prepared as described in Example 3), purified PPMT (prepared as described in Example 1), 50 mM MOPS buffer pH 7.2, 5 mM DTT, 1 mM EDTA, 1mg/ml BSA and 10 μl 3H-SAM were incubated at 37°C for 30 min and loaded onto a desalting column pre-equilibrated in 50 mM MOPS-Na pH 7.2, 1 mM DTT, 1 mM EDTA, 200 mM NaCl and 5% glycerol. The sample was eluted with the same buffer, 50 μl fractions were collected, and each aliquot analyzed by scintillation counting. Fractions containing ^3H -methyl PP2A were collected and stored at 4°C.

[00123] The following assay (see Lee, J. et al., Proc. Nat'l Acad. Sci. U.S. A. 93(12): 603-47 (1996)) is referred to subsequently as the "filter paper method". The test compound was incubated with purified PPME (prepared as described in Example 2) and purified ^3H -methyl-labeled PP2A (the preparation of which is described in the preceding paragraph) in Buffer C

(50 mM MOPS, 1mM EDTA, 1 mM DTT buffer at pH 7.2, as in Example 3). After incubation at 37°C for a fixed time (typically 10 minutes), the reaction mixture was spotted onto a 1 cm² piece of Whatman 3MM filter paper. The filter paper was immediately dipped in 10% ice-cold TCA, and then kept on ice in a plastic tray. After all reactions were run, the filter papers were washed at 4 °C with stirring in a large excess of 10% TCA, similarly washed twice at 4 °C with 100% methanol, and then vacuum dried at 45°C for 1 hr to remove the ³H-methanol produced by PP2A demethylation. The dried filter papers then were assayed for radioactivity by submerging them in scintillation fluid and performing scintillation counting. This procedure is easily adapted to a 96-well format to allow high-throughput screening (described below).

[00124] As used herein to refer to assay results, the term “inhibition activity” refers to the following relationship:

$$\text{Inhibition activity} = 1 - \frac{(\text{amount of methyl-PP2A demethylated with inhibitor})}{(\text{amount of methyl-PP2A demethylated without inhibitor})}$$

[00125] The activity of an inhibitor according to the present invention is defined herein as 1 unit of inhibitor inhibits 50% of the demethylation in 10 µl of reaction mixture in 10 minutes.

[00126] Drip-brewed coffee (prepared from Sumatra coffee purchased from Starbucks Coffee) was tested in an initial screen. The filter paper assay described above was used to assay 1 µl of Sumatra coffee per 10 µl of assay mixture. The assay showed that Sumatra coffee contained an inhibition activity of 0.83 (i.e., 83% inhibition relative to control).

[00127] Based on this significant inhibition of PPME, we tested an ethanol extract of Sumatra coffee that showed an even more significant inhibitory effect. 10g of ground Sumatra coffee bean (Starbucks) was extracted in ethanol at 80°C for 5 minutes with stirring. After cooling, the extract was filtered through a Whatman No. 54 paper filter. The filtered extract was rotary evaporated at 40°C until the volume was reduced to approximately 1 ml. The liquid phase was recovered and diluted into 50 mM MOPS buffer, 1 mM EDTA, pH 7.2.

[00128] Figure 3 shows the results of an assay using the filter paper method to assay 1µl of extract per 10 µl of assay mixture. The results show that this ethanol extract of Sumatra coffee grounds shows a concentration-dependent inhibition activity. The inhibitor effect seen is due to a small molecule that is soluble in organic solvents and is not caffeine. The

Concentration of pure caffeine required for 50% inhibition of PPME was $> 440 \mu\text{M}$. Inhibitory activity was present at similar levels in both regular and decaffeinated coffee.

Example 5: Assay for Methyltransferase activity

[00129] This example provides guidance for determining activation of PPMT. Purified PP2A AC dimers (see Example 3) are mixed with the purified PPMT (prepared as described in Example 1) and ^3H -methyl-labeled S-adenosyl methionine and incubated at 37°C . After several minutes aliquots were removed, protein was precipitated with 10% trichloroacetic acid (TCA), washed with 10% TCA, then washed with 100% methanol, and ^3H -methyl-PP2A was assayed by liquid scintillation counting (Tolstykh, T. et al., *EMBO J.* 19 (21): 5682-91 (2000)).

Example 6. Purification of the component(s) in coffee that inhibit PP2A MEase:

[00130] This example provides methods by which inhibition activity can be measured for a larger number of compounds in parallel in a 96-well plate format, and provides methods through which the inhibitory component(s) in coffee may be selectively enriched or purified.

(a) **Inhibitory effect of coffee component(s) on PPME assayed in a 96-well format:**

[00131] The inhibitory effect of coffee component(s) on PPME was measured using the filter paper method as adapted to a 96-well plate format. Equimolar amounts of PP2A and [^3H]-SAM and a two-fold molar excess of PPMT were incubated at 37°C for 30 min., which is the time when the methylation reaction reaches its equilibrium, and PP2A is close to 100% methylated. Methylated PP2A was separated from PPMT by diluting the reaction mixture in buffer with Ni-NTA resin and bovine serum albumin (BSA) and shaking at room temperature for 30 min. The supernatant containing methylated PP2A was collected, and Ni-resin to which PPMT bound was discarded.

[00132] 96-well Millipore filter plates were used to assay demethylation of methylated PP2A. Methylated PP2A was mixed with PPME at 37°C . At reaction times of 5 and 10 min, 10 μl of reaction mixture was spotted into a well of the filter plate containing cold 25% TCA to stop the demethylating reaction. The filter in each well was washed with 5% TCA, then with 70% ethyl alcohol, and finally air-dried. Scintillation fluid was added into the wells, and the radioactivity due to bound ^3H -methyl-PP2A was counted using a Packard TopCount scintillation counter.

(b) **Purification of coffee components inhibitory to PPME using ethyl acetate partitioning.**

[00133] The inhibition activity in coffee may be purified from extracts of either whole coffee or instant coffee by a selective partitioning into ethyl acetate at low versus high pH.

[00134] 10 g of ground Sumatra coffee bean (Starbucks) was extracted with ethanol at 80°C for 5 minute with stirring. After cooling, the extract was filtered through a Whatman No. 54 paper filter. The filtered extract was rotary evaporated at 40 °C until the volume was reduced to approximately 1 ml. 200 µl of this concentrated extract was mixed with an equal volume of 50 mM MOPS buffer, pH 7.2 with 1 mM EDTA (Buffer A), 0.1M HCl, or 0.1M NaOH, and then extracted with 1 ml of ethyl acetate. After separation by centrifugation, the aqueous and organic phases in each tube were collected, dried under vacuum in a Speedvac concentrator, resuspended in Buffer A to a volume of 2 ml, and the pH adjusted to 7.2. This extract was assayed using the filter paper method with a volume of 1 µl of extract per 10 µl of sample volume. The results of this assay demonstrated an inhibition activity of about 1 (100% inhibition) for the organic phase in the presence of Buffer A or HCl, an inhibition activity of about 0.2 for the organic phase in the presence of NaOH, an inhibition activity of about 0 from the aqueous phase in the presence of Buffer A or HCl, and an inhibition activity of about 1 from the aqueous phase in the presence of NaOH.

[00135] 1 g of decaffeinated instant coffee also was dissolved in a 10 ml final volume of 0.1M HCl (“acidic coffee solution”). The final pH of the acidic coffee solution was 3.0, and an amount of insoluble material was removed by centrifugation. The amount of insoluble material was larger than the amount of insoluble material present after dissolving 1 g of instant coffee in 10 ml of hot deionized water at pH 5 (the “water-extracted coffee solution”). 5 ml of the acidic coffee solution was extracted with 4 ml of ethyl acetate, and the organic phase recovered. The organic phase then was extracted with an equal volume of 0.1M NaOH. The final extract was clearer and much lighter in color than was instant coffee dissolved in water or HCl. These extracts were assayed using the 96 well plate method and a diluted extract equivalent to a volume of about 0.1 µl of extract per 10 µl of assay mixture. Under these conditions, the water extracted coffee solution had an inhibition activity of about 0.78, the acidic coffee solution had an inhibition activity of 0.44, and the NaOH extract of the organic phase had an inhibition activity of 0.47. These results and the results described above demonstrate that the inhibitory compounds from coffee can be purified by selective

partitioning into ethyl acetate at acidic pH and selective partitioning into the aqueous phase at basic pH.

(c) **Purification of coffee components inhibitory to PPME using ethanol extraction, water wash and reverse-phase separation.**

[00136] 10 g of decaffeinated instant coffee (Taster's Choice) was boiled (80°C) three times in 100 ml 100% ethanol for 5 min. Each ethanol extract was centrifuged at 3,000 rpm, 4°C for 30 min, the supernatant collected, and the coffee-solids-containing pellet used for the subsequent extraction. Inhibition activity present in these extracts was measured by the 96 well plate assay described in (a) above.

[00137] Figure 4A shows the presence of a methylation modifying compound possessing PP2A methylesterase inhibition activity in an extract prepared from instant coffee by ethanol extraction. Fig. 4A shows PPME inhibition activity in each crude ethanol extract after three 100 ml ethanol extractions, 300 ml total volume. The extracted inhibition activities in the second and third extracts were less than in the first extract, such that activity present in the second and third extractions was about half of the previous extract's activity. Based on this observation, 200ml EtOH should be enough to extract the majority of the methylation modifying activity in 10g of instant coffee.

[00138] Instant coffee was boiled (80°C) once in 100 ml ethanol for 5 minutes and twice time in 50ml ethanol for 5min each time. Each extract was centrifuged at 3,000 rpm, 4°C for 30 min, and the supernatants combined to yield a crude extract with an approximate volume of 200 ml. The inhibition activity in the final extract was measured by the 96 well plate assay described above.

[00139] Fig. 4B shows PP2A methyl esterase inhibition activity in crude ethanol extract after these extractions. As shown in Fig. 4B, about 10^6 units total inhibitory units were extracted in 200ml EtOH from 10g coffee.

[00140] The ethanol was filtered through Fisher Scientific medium porosity type P5 filter paper (P5 filter paper), rotary-evaporated and the brown, thick residue collected. The residue weighed 1.18 g. The dried coffee residue obtained from the crude ethanol extract was washed four times with 50 ml of doubly deionized water with pH < 5. The water solutions were filtered through P5 filter paper. 50ml of ethanol was used to dissolve components that stuck to the flask, and another 20ml ethanol was used to wash the filter paper. The material

dissolved in ethanol and not dissolvable in water was much lighter in color than was the crude extract.

[00141] The PP2A demethylation inhibition activity of the water washes, ethanol used to wash the flask, and ethanol used to wash the filter paper are shown in Fig. 4C. Only about 10% of the total inhibition activity in the crude ethanol extract was lost in the water-wash step. The residue dissolved from the flask and filter paper with ethanol was rotary evaporated to yield a final coffee extract containing nearly all the inhibitory activity and having a mass of about 0.1 g.

[00142] A solid-phase extraction cartridge containing C-18 resin (Waters) equilibrated with 50% methanol plus 0.1% trifluoroacetic acid (TFA) was used to further purify the inhibitory compound in the final coffee extract. The ethanol in the final coffee extract was rotary-evaporated. The residue was dissolved in 50% methanol plus 0.1% TFA and loaded on the C-18 cartridge. The cartridge then was washed with three column volumes of 50% methanol, 60% methanol, 70% methanol, 80% methanol, 90% methanol, 100% methanol, and 100% acetonitrile, all with 0.1% TFA. Materials eluted with each mobile phase were collected as separate fractions. Water and organic solvents in each fraction were rotary evaporated, and the residue was dissolved in ethanol.

[00143] Fig .4D shows the elution profile of inhibitory activity. Eluent corresponding to 0.5 µl of the final coffee extract was used per 10 µl of PP2A demethylation reaction. The 90% methanol and 100% methanol fractions contained most of the PPME inhibition activity present in the final coffee extract and little or no activity remained to be eluted by the acetonitrile.

Example 7. Identification of PP2A inhibitory activity in botanicals

[00144] We have used the assay described above in Example 4 to screen commercially available herbal substances for the presence of a PP2A demethylation inhibitor compound. Each of these substances has been reported to have properties that affect the health and well-being of human subjects. For example,

[00145] Allicin is an antibiotic and antifungal compound obtained from garlic.

[00146] Artichoke powder is believed to have hepatoprotective, antioxidant, and hypocholesterolemic properties.

[00147] Bacopa monniera is an herb used in India for memory, epilepsy, and as a mild sedative. Bacopa commonly grows in marshy areas throughout India. Bacopa is believed to have strong antioxidant properties, protect mental function and improve learning skills.

[00148] Butcher's broom, an herbal extract that typically contains 10% saponin glycosides as active components, has been reported to have anti-inflammatory properties.

[00149] Cocoa powder cocoa contains a high content of phenolic phytochemicals, or flavonoids, indicating the presence of known antioxidants that may protect against cancer, heart disease and other ailments.

[00150] Flaxseed oil, which is obtained by extracting the oil from flaxseeds is believed to stimulate the immune system and to have antioxidant, cholesterol and triglyceride lowering and anti-tumor properties.

[00151] Grape seed extract contains oligomeric proanthocyanidins (OPCs), a class of flavonoid complexes that act as antioxidants in the body. It is believed that OPCs may help protect against the effects of internal and environmental stresses and may counter the negative effects of high cholesterol on the heart and blood vessels.

[00152] Purportedly, nutritional supplementation of inositol may affect behavior and may have anti-depressant and anti-anxiety activities.

[00153] St. John's Wort, *Hypericum perforatum*, is an herbal product sold as an over-the-counter treatment for depression. It is being studied for its ability to lessen certain side effects of cancer treatment.

[00154] The antioxidants in green tea, black tea and red tea purportedly have significant health benefits, including the ability to prevent cancer and heart disease.

[00155] Tobacco extract has been shown to produce a biphasic effect on macrophage respiration: a stimulation at low concentrations and an inhibition at higher concentrations.

[00156] The putative health affects of ginseng, ginkgo, huperzine, Vitamin E (DL-alpha-tocopherol), and choline already have been described above.

[00157] Extracts typically were made from the contents of one tablet, capsule or caplet by heating in 1 ml of pure (200 proof) ethanol for 30 minutes at 80°C before centrifugation for 5 to 10 minutes at 12,000 rpm in a Sorvall MC-12V centrifuge to remove insoluble material. In the table below, this is indicated as a "standard ethanol" extraction. Extracts for tea were made by heating in 10 ml of doubly deionized water at 80°C for 30 minutes before centrifugation. This is denoted in the table as a "standard water" extraction. Other

extractions are as specified, and pure or nearly pure compounds were dissolved in the solvent listed. A portion of each extract was dried to determine the mass of dissolved material in the extract.

Table 1. Commercially available herbal substances screened for the presence of a PP2A demethylation inhibitor compound.

Sample Code	Substance	Extraction method	Source
#3	Gingko Biloba leaf extract: 60 mg per tablet. 24% flavonol glycosides (14.4 mg), 6% terpene lactones (3.6 mg). Also contains 14 support ingredients.	Standard ethanol	Sundown
#4	Allicin, (garlic bulb extract): 130 mg/capsule. Also contains cellulose.	Standard ethanol	Zhang
#5	Huperzine A, Gotu Kola (Centella asiatica, aerial part) 285 mg, Huperzine 50µg, lecithin 100 mg/capsule. Also contains 3 support ingredients.	Dissolved in ethanol	Solaray
#6	Bacopa, Himalaya (Bacopa moniera fennel leaf extract): 500 mg/capsule.	Standard ethanol	Brahmi
#7	Butcher's Broom: root extract with other ingredients, no amount was shown on the bottle label. Also it is not clear whether or not support ingredient was included.	Standard ethanol	TWINLAB Nature's Herbs
#10	Flaxseed oil - contains 1425 mg of linolenic acid, 450 mg of oleic acid, 400 mg of linoleic acid, 150 mg of palmitic acid, 75 mg of stearic acid/2 softgels. Other ingredients: flaxseed oil. Oil volume is about 1.4 ml/soft gel.	Dissolved in DMSO	Solgar
#12-1	DL- α -tocopherol, , 95%, MW 431, prepared from synthetic phytol.	Dissolved in ethanol	Sigma (T3251-25G)
#21	Ginseng - 100 mg of Ginseng extract / caplet, 7% ginsenosides in the extract, also contains calcium carbonate et al 16	Standard ethanol	Your Life

Sample Code	Substance	Extraction method	Source
	ingredients		
#22	Grape Seed Extract, 100 mg of grape seed extract (85% polyphenols) / tablet, also contains calcium carbonate et al 11 ingredients.	Standard ethanol	Wegmans
#23	St. John's Wort, 300 mg of St. John's wort extract (0.3 % total dianthrones) / caplet, , also contains cellulose et al 11 ingredients.	Standard ethanol	Wegman's
#24	Artichoke-powder, 100 mg of artichoke (aerial part) extract and 375 mg of artichoke (aerial part) /capsule, also contains vitamin E et al 4 ingredients.	Standard ethanol	Nature's Herbs
#25	Choline & Inositol Capsules, choline 250 mg and inositol 250 mg/ capsule, also contains gelatin et al 5 ingredients.	Extracted with room temperature ethanol; otherwise standard.	Twinlab
#26	Green Tea, 1.5 g/bag, ingredients: green tea, verbena, lemongrass, lemon flavor.	Standard water	Wissotzky
#27	Red Bush Tea, 2 g/bag, Ingredients: South African rooibos (<i>Aspalathus linearis</i>) and lemongrass with 15 mg of SGS made from broccoli seed. (no caffeine)	Standard water	Brassica
#28	Black Tea, 2.1 g/bag, Chinese black tea with 15 mg of SGS made from broccoli seed, caffeine	Standard water	Brassica
#30E	Tobacco No additive. Sliced tobacco leaves.	820 mg extracted with 10 ml ethanol for 60 min at 80°C	Natural American Spirit
#31	Cocoa, unsweetened -.	150 mg of the powder was extracted using standard ethanol extraction.	Ghiradelli Chocolate Co.
CCM	Curcumin	Dissolved in ethanol.	LKT Laboratories, Inc.
CGA	Chlorogenic acid ("CGA"), predominantly trans(ex coffee seeds) 99%,MW=354.3	Dissolved in 50% ethanol.	Acros Organics

Sample Code	Substance	Extraction method	Source
DHCA	3,4-Dihydroxycinnamic acid ("DHCA"), predominantly trans isomer 99+%, MW=180.6	Dissolved in 50% ethanol.	Acros Organics
HMCA	4-Hydroxy-3-methoxycinnamic acid ("HMCA"), 99%, MW=194	Dissolved in 50% ethanol.	Acros Organics
QA	D(-)-Quinic acid ("QA"), 98%, MW192.17,	Dissolved in buffer.	Acros Organics

[00158] The concentration of each substance that resulted in an inhibition activity of 0.5 (50% inhibition of PPME) was determined using the filter paper method described above in Example 4, and is hereafter referred to as the IC₅₀. The results, which are expressed as dry extract weight/ml except as noted, are shown in Table 2.

Table 2. Effect of Substances on PPME activity (assayed against PPME at 85 nM by the filter paper method.

Sample Code	Substance name	IC ₅₀
# 3	Ginkgo (extract)	106 µg/ml
# 4	Allicin (extract)	100 µg/ml
# 5-1	Huperzine	No inhibition at 40 µM
# 6	Bacopa (extract)	63 µg/ml
# 7	Butcher's broom	37 µg/ml
# 10	Flaxseed oil	1.3 mg/ml
#12-1	DL-α-tocopherol	35 µM
# 21	Ginseng (extract)	360 µg/ml
# 22	Grape seed (extract)	25 µg/ml
# 23	St. John's wort (extract)	22 µg/ml
# 24	Artichoke-powder (extract)	260 µg/ml
# 25	Choline & inositol caps (extract)	75 µg/ml
# 26	Green tea (extract)	179 µg/ml
# 27	Red bush tea (extract)	45 µg/ml
# 28	Black tea (extract)	50 µg/ml
# 30E	Tobacco (extract)	344 µg/ml
# 31	Cocoa, Ghirardelli Chocolate Co., unsweetened	92 µg/ml
CCM-L	Curcumin-(LKT Co.)	771 µM
CGA	Chlorogenic acid	> 1 mM
DHCA	Dihydroxycinnamic acid	> 1 mM
HMCA	Hydromethoxycinnamic acid	>1 mM
QA	Quinic acid	> 10 mM

[00159] For purposes of this table, the unit of concentration in the above table ($\mu\text{g/ml}$ or mg/ml) refers to dried extract weight (DEW) unless otherwise noted. A “>” symbol before the IC_{50} value means that the IC_{50} concentration is higher than the value tested. Most of these IC_{50} values are high compared to the concentration of the AC dimer of PP2A (30-50 nM) in the reaction mixture.

[00160] From these results we conclude that all of these compounds except huperzine, chlorogenic acid, dihydroxycinnamic acid, HMCA, and quinic acid contain significant PPME inhibitory activity. Since the measured activity is dependent on the extraction procedure, the assay described can be used to optimize the extraction protocol and to select varieties or sources of these compounds that contain the most concentrated PPME inhibitory activity.

[00161] The present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the Invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the following claims.

CLAIMS

What is claimed is:

1. A composition for promoting general health and well-being in a mammalian subject, including a human, comprising an isolated methylation modifying compound.
2. The composition according to claim 1, wherein the compound is isolated from a botanical raw material.
3. The composition according to claim 1, wherein the compound is isolated from a microbial raw material.
4. The composition according to claim 1, wherein the composition inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme.
5. The composition according to claim 4, wherein the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme.
6. The composition according to claim 4, wherein the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase.
7. The composition according to claim 6, wherein the demethylating activity of the protein phosphatase 2A specific protein methylesterase is determined by measuring levels of protein phosphatase 2A methyl esterification.
8. The composition according to claim 2, wherein the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; coffee; tea; tobacco; and cocoa.
9. The composition according to claim 2, wherein the botanical raw material comprises a fruit of a species of plant genus *Coffea*.
10. The composition according to claim 9, wherein the fruit of the plant genus *Coffea* is a coffee bean.
11. The composition according to claim 2, wherein the botanical raw material comprises a nutritional supplement.
12. The composition according to claim 11, wherein the nutritional supplement comprises at least one tocopherol.

13. The composition according to claim 11, wherein the nutritional supplement comprises Vitamin E.
14. The composition according to claim 2, wherein the botanical raw material comprises a leaf of a tobacco plant.
15. The composition according to claim 2, wherein the botanical raw material comprises a ground coffee.
16. The composition according to claim 2, wherein the botanical raw material comprises an instant coffee.
17. The composition according to claim 2, wherein the methylation modifying compound is isolated from the botanical raw material by extracting the botanical raw material with a polar solvent.
18. The composition according to claim 17, wherein the polar solvent is a lower alcohol.
19. The composition according to claim 18, wherein the lower alcohol is ethanol.
20. The composition according to claim 17, wherein the polar solvent is ethyl acetate.
21. The composition according to claim 20, wherein the methylation modifying compound is soluble in ethyl acetate below a pH of about 5 and insoluble in ethyl acetate above a pH of about 10.
22. The composition according to claim 1, wherein the composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.
23. A comestible for promoting general health and well-being in a mammalian subject, including a human, comprising a composition containing an isolated methylation modifying compound.
24. The comestible according to claim 23, wherein the compound is isolated from a botanical raw material.
25. The comestible according to claim 23, wherein the compound is isolated from a microbial raw material.

26. The comestible according to claim 23, wherein the composition inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme.

27. The comestible according to claim 26, wherein the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme.

28. The comestible according to claim 26, wherein the demethylating enzyme is protein phosphatase 2A specific protein methylesterase.

29. The comestible according to claim 26, wherein the level of protein methylesterase demethylating activity is determined by measuring levels of protein phosphatase 2A methyl esterification.

30. The comestible according to claim 24, wherein the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; coffee; tea; tobacco; and cocoa.

31. The comestible according to claim 24, wherein the botanical raw material comprises a fruit of a species of plant genus *Coffea*.

32. The comestible according to claim 31, wherein the fruit of the plant genus *Coffea* is a coffee bean.

33. The comestible according to claim 24, wherein the botanical raw material comprises a nutritional supplement.

34. The comestible according to claim 33, wherein the nutritional supplement comprises at least one tocopherol.

35. The comestible according to claim 33, wherein the nutritional supplement comprises Vitamin E.

36. The comestible according to claim 24, wherein the botanical raw material comprises a leaf of a tobacco plant.

37. The comestible according to claim 24, wherein the botanical raw material comprises a ground coffee.

38. The comestible according to claim 24, wherein the botanical raw material comprises an instant coffee.

39. The comestible according to claim 24, wherein the methylation modifying compound is isolated from the botanical raw material by extracting the botanical raw material with a polar solvent.

40. The comestible according to claim 39, wherein the polar solvent is a lower alcohol.

41. The comestible according to claim 40, wherein the lower alcohol is ethanol.

42. The comestible according to claim 39, wherein the polar solvent is ethyl acetate.

43. The comestible according to claim 42, wherein the methylation modifying compound is soluble in ethyl acetate below a pH of about 5 and insoluble in ethyl acetate above a pH of about 10.

44. The comestible according to claim 23, wherein the composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.

45. The comestible according to claim 23, wherein the comestible is a beverage.

46. The comestible according to claim 45, wherein the beverage is selected from the group consisting of a drink comprising water, a fruit drink, a coffee, a tea, an energy drink, a baby formula, an adult nutritional drink, a health drink, and a sports drink.

47. The comestible according to claim 23, wherein the comestible is a food.

48. The comestible according to claim 47, wherein the comestible is a cereal.

49. The comestible according to claim 23, wherein the comestible is a chewing gum.

50. The comestible according to claim 23, wherein the comestible is a candy.

51. The comestible according to claim 23, wherein the comestible is an ingredient of an inhalant.

52. The comestible according to claim 23, wherein the comestible is an ingredient of a transdermal delivery system.

53. A pharmaceutical preparation for promoting general health and well-being in a mammalian subject, including a human, comprising a cognitive function-enhancing amount

of a composition containing an isolated methylation modifying compound and a pharmaceutically acceptable carrier.

54. The pharmaceutical preparation according to claim 53, wherein the compound is isolated from a botanical raw material.

55. The pharmaceutical preparation according to claim 53, wherein the compound is isolated from microbial raw material.

56. The pharmaceutical preparation according to claim 53, wherein the compound inhibits a demethylating enzyme that acts on a protein phosphatase 2A and thereby stimulates methylation of the protein phosphatase 2A enzyme.

57. The pharmaceutical preparation according to claim 56, wherein the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme.

58. The pharmaceutical preparation according to claim 56, wherein the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase.

59. The pharmaceutical preparation according to claim 58, wherein the demethylating activity of the protein phosphatase 2A specific protein methylesterase is determined by measuring levels of protein phosphatase 2A methyl esterification.

60. The composition according to claim 54, wherein the botanical raw material comprises at least one substance selected from the group consisting of ginkgo; allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline, inositol; tea; tobacco; and cocoa.

61. The pharmaceutical preparation according to claim 54, wherein the botanical raw material comprises a fruit of a species of plant genus *Coffea*.

62. The pharmaceutical preparation according to claim 61, wherein the fruit of the plant genus *Coffea* is a coffee bean.

63. The pharmaceutical preparation according to claim 54, wherein the botanical raw material comprises a ground coffee.

64. The pharmaceutical preparation according to claim 54, wherein the botanical raw material comprises an instant coffee.

65. The pharmaceutical preparation according to claim 54, wherein the botanical raw material is a nutritional supplement.

REPLACEMENT PAGES

66. The pharmaceutical preparation according to claim 65, wherein the nutritional supplement comprises at least one tocopherol.

67. The pharmaceutical preparation according to claim 65, wherein the nutritional supplement comprises vitamin E.

68. The pharmaceutical preparation according to claim 54, wherein the botanical raw material comprises a leaf of a tobacco plant.

69. The pharmaceutical preparation according to claim 54, wherein the methylation modifying compound is isolated from the botanical raw material by extracting the botanical raw material with a polar solvent.

70. The pharmaceutical preparation according to claim 69, wherein the polar solvent is a lower alcohol.

71. The pharmaceutical preparation according to claim 70, wherein the lower alcohol is ethanol.

72. The pharmaceutical preparation according to claim 69, wherein the polar solvent is ethyl acetate.

73. The pharmaceutical preparation according to claim 72, wherein the methylation modifying compound is soluble in ethyl acetate below a pH of about 5 and insoluble in ethyl acetate above a pH of about 10.

74. The pharmaceutical preparation according to claim 53, wherein the preparation is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.

75. The pharmaceutical preparation according to claim 53, wherein the preparation is administered orally.

76. The pharmaceutical preparation according to claim 53, wherein the preparation is in the form of a powder.

77. The pharmaceutical preparation according to claim 53, wherein the preparation is in the form of a tablet.

78. The pharmaceutical preparation according to claim 53, wherein the preparation is in the form of a capsule.

79. The pharmaceutical preparation according to claim 53, wherein the preparation is in the form of a lozenge.

80. The pharmaceutical preparation according to claim 53, wherein the preparation is in the form of a suppository.

81. A method for promoting general health and well-being in a mammalian subject, including a human, the method comprising the steps

(a) isolating a methylation modifying compound; and

(b) administering a cognitive function-enhancing amount of a composition comprising the isolated methylation modifying compound to a subject in need thereof.

82. The method according to claim 80, wherein in step (a) the methylation modifying compound is isolated from a botanical raw material.

83. The method according to claim 80, wherein in step (a) the methylation modifying compound is isolated from a microbial raw material.

84. The method according to claim 80, wherein the methylation modifying compound inhibits a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme.

85. The method according to claim 83, wherein the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme.

86. The method according to claim 83, wherein the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase.

87. The method according to claim 85, further comprising the step of (c) determining the demethylating activity of the protein phosphatase 2A specific protein methylesterase.

88. The method according to claim 80, wherein the cognition function-enhancing amount of the composition is from about 1 mg to about 10 g.

89. The method according to claim 81, wherein the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin, bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; tea; tobacco; and cocoa.

90. The method according to claim 81, wherein the botanical raw material comprises a fruit of a species of plant genus *Coffea*.

91. The method according to claim 89, wherein the fruit of the plant genus *Coffea* is a coffee bean.

92. The method according to claim 81, wherein the botanical raw material comprises a ground coffee.

93. The method according to claim 81, wherein the botanical raw material comprises an instant coffee.

94. The method according to claim 81, wherein the botanical raw material comprises a nutritional supplement.

95. The method according to claim 93, wherein the nutritional supplement comprises at least one tocopherol.

96. The method according to claim 93, wherein the nutritional supplement comprises vitamin E.

97. The method according to claim 81, wherein the botanical raw material comprises a leaf of a tobacco plant.

98. The method according to claim 81, step (a) further comprising the step of extracting the botanical raw material with a polar solvent.

99. The method according to claim 97, wherein the polar solvent is a lower alcohol.

100. The method according to claim 98, wherein the lower alcohol is ethanol.

101. The method according to claim 97, wherein the polar solvent is ethyl acetate.

102. The method according to claim 100, step (a) further comprising the steps of solubilizing the methylation modifying compound in ethyl acetate below a pH of about 5 and partitioning the methylation modifying compound between an aqueous phase and ethyl acetate above a pH of about 10.

103. The method according to claim 80, wherein the extracted composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.

104. The method according to claim 80, wherein in step (b), the composition is administered orally.

105. The method according to claim 80, wherein the composition is a powder.

106. The method according to claim 80, wherein the composition is in the form of a tablet

107. The method according to claim 80, wherein the composition is in the form of a capsule.

108. The method according to claim 80, wherein the composition is in the form of a lozenge.

109. The method according to claim 80, wherein the composition is in the form of a suppository.

110. A composition for promoting general health and well-being in a mammalian subject, including a human, prepared by a process, the process comprising the steps of:

a) contacting a botanical raw material with a polar solvent for a time sufficient to form an extract solution from the botanical raw material;

b) isolating the extract solution and removing the polar organic solvent to form a concentrated primary extract; and

c) dissolving the composition in a solvent at a pH ranging from about 7 to about 10 to form a concentrated purified extract.

111. The composition according to claim 109 wherein the polar solvent in step (a) of the process is a lower alcohol.

112. The composition according to claim 110, wherein the lower alcohol is ethanol.

113. The composition according to claim 119, wherein the polar solvent in step (a) of the process is ethyl acetate.

114. The composition according to claim 112, wherein step (a) of the process further comprises the step of solubilizing the composition in ethyl acetate below a pH of about 5.

115. The composition according to claim 112, wherein step (b) of the process further comprises the step of partitioning the methylation modifying compound between an aqueous phase and ethyl acetate above a pH of about 10.

116. The composition according to claim 109, wherein the composition inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme.

117. The composition according to claim 115, wherein the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme.

118. The composition according to claim 115, wherein the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase.

119. The composition according to claim 109, wherein the botanical raw material comprises at least one substance selected from the group consisting of ginkgo; allicin; bacopa; butcher's broom; flaxseed oil; a tocopherol; vitamin E; ginseng; grape seed,; St. John's wort; artichoke-powder; choline; inositol; coffee; tea; tobacco; and cocoa.

120. The composition according to claim 109, wherein the botanical raw material comprises a fruit of a species of plant genus *Coffea*.

121. The composition according to claim 119, wherein the fruit of the plant genus *Coffea* is a coffee bean.

122. The composition according to claim 109, wherein the botanical raw material comprises a ground coffee.

123. The composition according to claim 109, wherein the botanical raw material comprises an instant coffee.

124. The composition according to claim 109, wherein the botanical raw material comprises a nutritional supplement.

125. The composition according to claim 123, wherein the nutritional supplement comprises at least one tocopherol.

126. The composition according to claim 123, wherein the nutritional supplement comprises Vitamin E.

127. The composition according to claim 109, wherein the botanical raw material comprises a leaf of a tobacco plant.

128. The composition according to claim 109, wherein the concentrated purified extract is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.

129. The composition according to claim 109, wherein the composition is administered orally.

130. The composition according to claim 109, wherein the composition is in the form of a powder.

131. The composition according to claim 109, wherein the composition is in the form of a tablet.

132. The composition according to claim 109, wherein the composition is in the form of a capsule.

133. The composition according to claim 109, wherein the composition is in the form of a lozenge.

134. The composition according to claim 109, wherein the composition is in the form of a suppository.

135. A method of treating or preventing cognitive effects of a disease, condition, or disorder involving defective methylation metabolism in a mammalian subject, including a human, the method comprising the steps of

- (a) isolating a methylation modifying compound; and
- (b) administering a therapeutically effective amount of a composition comprising the isolated methylation modifying compound to a subject in need thereof.

136. The method according to claim 134, wherein in step (a) the methylation modifying compound is isolated from a botanical raw material.

137. The method according to claim 134, wherein in step (a) the methylation modifying compound is isolated from a microbial raw material.

138. The method according to claim 134, wherein the disease, condition, or disorder is associated with higher than normal levels of plasma homocysteine.

139. The method according to claim 134, wherein the disease condition or disorder is at least one disease, condition or disorder selected from the group consisting of a cardiac disorder; atherosclerosis; a neurodegenerative disorder; a cerebrovascular disorder; a neuropsychiatric disorder; and diabetes.

140. The method according to claim 134, wherein the methylation modifying compound inhibits a demethylating activity of a demethylating enzyme that acts on a protein phosphatase 2A enzyme and thereby stimulates methylation of the protein phosphatase 2A enzyme.

141. The method according to claim 139, wherein the composition inhibits at least about 50% of the demethylating activity of the demethylating enzyme.

142. The method according to claim 139, wherein the demethylating enzyme is a protein phosphatase 2A specific protein methylesterase.

143. The method according to claim 134, wherein the therapeutically effective amount of the composition is from about 1 mg to about 10 g of the composition.

144. The method according to claim 134, wherein the botanical raw material comprises at least one substance selected from the group consisting of ginkgo, allicin,

bacopa; butcher's broom; flaxseed oil; a tocopherol; Vitamin E; ginseng, grape seed, St. John's wort; artichoke-powder; choline; inositol; tea; tobacco; and cocoa.

145. The method according to claim 134, wherein the botanical raw material comprises a fruit of a species of plant genus *Coffea*.

146. The method according to claim 144, wherein the fruit of the plant genus *Coffea* is a coffee bean.

147. The method according to claim 134, wherein the botanical raw material comprises a ground coffee.

148. The method according to claim 134, wherein the botanical raw material comprises an instant coffee.

149. The method according to claim 134, wherein the botanical raw material comprises a nutritional supplement.

150. The method according to claim 148, wherein the nutritional supplement comprises at least one tocopherol.

151. The method according to claim 148, wherein the nutritional supplement comprises Vitamin E.

152. The method according to claim 134, wherein the botanical raw material comprises a leaf of a tobacco plant.

153. The method according to claim 134, step (a) further comprising the step of extracting the botanical raw material with a polar solvent.

154. The method according to claim 152, wherein the polar solvent is a lower alcohol.

155. The method according to claim 152, wherein the polar solvent is ethyl acetate.

156. The method according to claim 154, wherein step (a) further comprises the steps of solubilizing the methylation modifying compound in ethyl acetate below a pH of about 5 and partitioning the methylation modifying compound between an aqueous phase and ethyl acetate above a pH of about 10.

157. The method according to claim 134, wherein the composition is essentially free of at least one substance selected from the group consisting of caffeine, caffeic acid and chlorogenic acid.

158. The method according to claim 134, wherein in step (b) the composition is administered orally.

- 159. The method according to claim 134, wherein the composition is a powder.
- 160. The method according to claim 134, wherein the composition is in the form of a tablet.
- 161. The method according to claim 134, wherein the composition is in the form of a capsule.
- 162. The method according to claim 134, wherein the composition is in the form of a lozenge.
- 163. The method according to claim 134, wherein the composition is in the form of a suppository.

FIG. 1/7




FIG. 2/7





FIG. 3/7

FIG. 4A/7





FIG. 4B/7

FIG. 4D/7





FIG. 4C/7



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US06/03686

A. CLASSIFICATION OF SUBJECT MATTER

IPC: A61K 36/00(2006.01)

USPC: 424/776

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/776

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Searched West

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,502,056 A (BREITBARTH) 26 March 1996 (26.03.1996), column 2, lines 14-24.	-162
Y	US 6,399,796 B2 (SCHWATZ) 04 June 2002 (04.06.2002), column 5, lines 50-62.	1-162
Y	US 6,760,359 B1 (COPELAND et al.) 15 June 2004 (15.06.2004), column 2, lines 7-18.	1-162

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:		"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

31 May 2006 (31.05.2006)

Date of mailing of the international search report

15 JUN 2006

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (571) 273-3201

Authorized officer

Randall Winston

Telephone No. (571) 272-1600